

Journal of Hyperstructures 12 (1) (2023), 30-50.

<https://doi.org/10.22098/jhs.2023.2511>

Research Paper

FUZZY α -MODULARITY IN FUZZY α -LATTICES

Payal A. Khubchandani  and Jyoti A. Khubchandani

ABSTRACT. In this paper, we have introduced and studied the notion of a fuzzy independent pair and obtain some properties of fuzzy α -modular pairs and independent pairs.

Key Words: Fuzzy α -lattice, fuzzy α -modular pair, fuzzy atom, fuzzy independent pair, \perp_F -symmetric, fuzzy semi-modular.

2010 Mathematics Subject Classification: Primary: 06D72; Secondary: 03E72.

1. INTRODUCTION

Zadeh [14] in 1971 introduced the concept of fuzzy ordering. The concept of a fuzzy sublattice was introduced by Yuan and Wu [13]. Ajmal and Thomas [1] in 1994 defined a fuzzy lattice and a fuzzy sublattice as a fuzzy algebra. Chon [2] considered Zadeh's fuzzy order [15] and proposed a new notion of a fuzzy lattice and studied level sets of such structures. At the same time, he also proved some results for distributive and modular fuzzy lattices. Mezzomo *et. al.* [4] changed the way to define the fuzzy supremum and the fuzzy infimum of a pair of elements by considering a threshold fixed $\alpha \in [0, 1)$ instead of, as usual, zero.

The concept of a modular pair in a lattice is well investigated by Maeda and Maeda [3]. Wasadikar and Khubchandani [7] defined a fuzzy modular pair in a fuzzy lattice and obtained some properties of fuzzy

Received: 17 April 2023, Accepted: 30 June 2023. Communicated by Mahmood Bakhshi ;
*Address correspondence to P.A. Khubchandani; E-mail: payal.khubchandani@yahoo.com.

This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](#).

Copyright © 2023 The Author(s). Published by University of Mohaghegh Ardabili.

modular pairs. Recently, Wasadikar and Khubchandani [12] introduced the notion of a fuzzy α -modular pair in a fuzzy α -lattice and prove some properties analogous to classical theory. In this paper, we introduce and study the notion of a fuzzy independent pair and obtain some properties of fuzzy α -modular pairs and independent pairs in fuzzy α -lattice.

2. PRELIMINARIES

In fuzzy sets, each element of a nonempty set X is mapped to $[0, 1]$ by a membership function $\mu : X \rightarrow [0, 1]$.

A mapping $A : X \times X \rightarrow [0, 1]$ is called a fuzzy binary relation on X .

The following definition is from Zadeh [15]. A fuzzy binary relation A on X is called:

- (i) fuzzy reflexive if $A(x, x) = 1$, for all $x \in X$;
- (ii) fuzzy symmetric if $A(x, y) = A(y, x)$, for all $x, y \in X$;
- (iii) fuzzy transitive if $A(x, z) \geq \sup_{y \in X} \min[A(x, y), A(y, z)]$;
- (iv) fuzzy antisymmetric if $A(x, y) > 0$ and $A(y, x) > 0$ implies $x = y$.

Based on the above properties Zadeh [15] introduced the following concepts related to a fuzzy binary relation A on a set X :

- (i) A is called a fuzzy equivalence relation on X if A is fuzzy reflexive, fuzzy symmetric and fuzzy transitive;
- (ii) A is a fuzzy partial order relation if A is fuzzy reflexive, fuzzy antisymmetric and fuzzy transitive and the pair (X, A) is called a fuzzy partially ordered set or a fuzzy poset;
- (iii) A is a fuzzy total order relation if it is a fuzzy partial order relation and $A(x, y) > 0$ or $A(y, x) > 0$, for all $x, y \in X$, and the fuzzy poset (X, A) is called of a fuzzy totally ordered set or a fuzzy chain.

Definition 2.1. [2, Definition 3.1] Let (X, A) be a fuzzy poset and let $Y \subseteq X$. An element $u \in X$ is said to be an upper bound for Y iff $A(y, u) > 0$, for all $y \in Y$. An upper bound u_0 for Y is the least upper bound (or supremum) of Y iff $A(u_0, u) > 0$, for every upper bound u for Y . We then write $u_0 = \sup Y = \vee Y$. If $Y = \{x, y\}$, then we write $\vee Y = x \vee y$.

Similarly, an element $v \in X$ is said to be a lower bound for Y iff $A(v, y) > 0$, for all $y \in Y$. A lower bound v_0 for Y is the greatest lower bound (or infimum) of Y iff $A(v, v_0) > 0$, for every lower bound v for

Y . We then write $v_0 = \inf Y = \wedge Y$. If $Y = \{x, y\}$, then we write $\wedge Y = x \wedge y$.

3. FUZZY α -LATTICES

Mezzomo and Bedregal [4] generalized the concept of a (fuzzy) upper bound as follows.

Definition 3.1. [4, Definition 3.1] Let (X, A) be a fuzzy poset. Let $Y \subseteq X$ and $\alpha \in [0, 1]$. An element $u \in X$ is said to be an α -upper bound for Y whenever $A(x, u) > \alpha$, for all $x \in Y$. An α -upper bound u_0 for Y is called a least α -upper bound (or α -Supremum) of Y iff $A(u_0, u) > \alpha$, for every α -upper bound u of Y .

Dually, an element $v \in X$ is said to be an α -lower bound for Y iff $A(v, x) > \alpha$, for all $y \in Y$. An α -lower bound v_0 for Y is called a greatest α -lower bound (or α -infimum) of Y iff $A(v, v_0) > \alpha$ for every α -lower bound v for Y .

We denote the least α -upper bound of the set $\{x, y\}$ by $x \vee_\alpha y$ and the greatest α -lower bound of the set $\{x, y\}$ by $x \wedge_\alpha y$.

Remark 3.2. [4, Remark 3.1] Since A is fuzzy antisymmetric, the least α -upper (greatest α -lower) bound, if it exists, is unique.

Proposition 3.3. [4, Proposition 3.1] Let (X, A) be fuzzy poset, $\alpha \in [0, 1)$ and $x, y, z \in X$. If $A(x, y) > \alpha$ and $A(y, z) > \alpha$, then $A(x, z) > \alpha$.

Definition 3.4. [4, Definition 3.2] A fuzzy poset (X, A) is a fuzzy α -lattice iff $x \vee_\alpha y$ and $x \wedge_\alpha y$ exists for all $x, y \in X$, for some $\alpha \in [0, 1)$.

Definition 3.5. [4, Definition 3.4] A fuzzy poset (X, A) is called fuzzy sup α -lattice, if each pair of element has α -supremum in X , denoted by $\sup_\alpha X$.

Dually, it is called fuzzy inf α -lattice, if each pair of element has α -infimum in X , denoted by $\inf_\alpha X$. A fuzzy semi α -lattice is a fuzzy poset which is a fuzzy sup α -lattice or a fuzzy inf α -lattice.

Definition 3.6. [4, Definition 3.5] Let (X, A) be a fuzzy poset and I be a fuzzy set on X . The α -supremum in I denoted by $\sup_\alpha I$, is an element of X such that if $x \in X$ and $\mu_I(x) > \alpha$, then $A(x, \sup_\alpha I) > \alpha$ and if $u \in X$ is such that $A(x, u) > \alpha$ whenever $\mu_I(x) > \alpha$, then $A(\sup_\alpha I, u) > \alpha$.

Similarly, the α -infimum in I denoted by $\inf_\alpha I$, is an element of X such that if $x \in X$ and $\mu_I(x) > \alpha$, then $A(\inf_\alpha I, x) > \alpha$ and if $v \in X$ is such that $A(v, x) > \alpha$ whenever $\mu_I(x) > \alpha$, then $A(v, \inf_\alpha I) > \alpha$.

Definition 3.7. [4, Definition 3.6] A fuzzy inf α -lattice is called inf complete if all of its nonempty fuzzy sets have α -infimum.

Similarly, a fuzzy sup α -lattice is called sup-complete if all of its nonempty fuzzy sets admit α -supremum. A fuzzy α -lattice is complete whenever it is, simultaneously, inf-complete and sup-complete.

Proposition 3.8. [4, Proposition 3.2] Let (X, A) be a complete fuzzy sup α -lattice (inf α -lattice) and I be a fuzzy set on X . Then, $\sup_\alpha I$ ($\inf_\alpha I$) exists and it is unique.

Proposition 3.9. [4, Proposition 3.3] Let $\mathcal{L} = (X, A)$ be a fuzzy sup α -lattice, then there exist an element \top in X , such that $A(x, \top) > \alpha$ for all $x \in X$.

Proposition 3.10. [4, Proposition 3.4] Let $\mathcal{L} = (X, A)$ be a fuzzy inf α -lattice, then there exist an element \perp in X , such that $A(\perp, x) > \alpha$ for all $x \in X$.

Definition 3.11. [4, Definition 3.6] A fuzzy lattice (X, A) is bounded if there exists \top and \perp in X such that for any $x \in X$, $A(\perp, x) > \alpha$ and $A(x, \top) > \alpha$.

Corollary 3.12. [4, Corollary 3.1] Every fuzzy lattice is a fuzzy α -lattice.

We illustrate the concepts of an α -upper bound and α -lower bound with an example.

Example 3.13. Consider the set $X = \{x, y, z, w\}$, let $\alpha=0.2$ and let $A : X \times X \longrightarrow [0, 1]$ be a fuzzy relation defined as follows:

$$A(x, x) = A(y, y) = A(z, z) = A(w, w) = 1.0,$$

$$A(w, z) = 0.2, A(w, y) = 0.5, A(w, x) = 0.9,$$

$$A(z, w) = 0.0, A(z, y) = 0.3, A(z, x) = 0.6,$$

$$A(y, w) = 0.0, A(y, z) = 0.0, A(y, x) = 0.4,$$

$$A(x, w) = 0.0, A(x, z) = 0.0, A(x, y) = 0.0.$$

Then A is a fuzzy total order relation.

Let $Y = \{w, z\}$. Then x, y and z are the α -upper bounds of Y . Since $A(z, w) = 0.0$ and $A(w, z) = 0.2 \geq \alpha$, it follows that the α -supremum of Y is z and the α -infimum is w .

The fuzzy α -join and fuzzy α -meet tables are as follows:

\vee_α	x	y	z	w	\wedge_α	x	y	z	w
x	x	x	x	x	x	x	y	z	w
y	x	y	y	y	y	y	y	z	w
z	x	y	z	z	z	z	z	z	w
w	x	y	z	w	w	w	w	w	w

We note that (X, A) is a fuzzy lattice as well as a fuzzy α -lattice for $\alpha = 0.2$.

In Figure 1, we show the related tabular and graphical representations for the fuzzy relation A .

A	w	z	y	x
w	1.0	0.2	0.5	0.9
z	0.0	1.0	0.3	0.6
y	0.0	0.0	1.0	0.4
x	0.0	0.0	0.0	1.0




Figure 1

The following example shows that a subset of a fuzzy poset may not have a greatest α -lower bound (least α -upper bound).

Example 3.14. Let $X = \{x_1, y_1, z_1, w_1\}$.

Let $A : X \times X \rightarrow [0, 1]$ be a fuzzy relation defined as follows:

$$A(x_1, x_1) = A(y_1, y_1) = A(z_1, z_1) = A(w_1, w_1) = 1.0,$$

$$A(x_1, y_1) = 0.20, A(x_1, z_1) = 0.30, A(x_1, w_1) = 0.90,$$

$$A(y_1, x_1) = 0.0, A(y_1, z_1) = 0.0, A(y_1, w_1) = 0.50,$$

$$A(z_1, x_1) = 0.0, A(z_1, y_1) = 0.0, A(z_1, w_1) = 0.70,$$

$$A(w_1, x_1) = 0.0, A(w_1, y_1) = 0.0, A(w_1, z_1) = 0.0.$$

Then A is a fuzzy partial order relation.

The fuzzy α -join and fuzzy α -meet tables are as follows:

\vee_α	x_1	y_1	z_1	w_1	\wedge_α	x_1	y_1	z_1	w_1
x_1	x_1	y_1	z_1	w_1	x_1	x_1	x_1	x_1	x_1
y_1	y_1	y_1	w_1	w_1	y_1	x_1	y_1	x_1	y_1
z_1	z_1	w_1	z_1	w_1	z_1	x_1	x_1	z_1	z_1
w_1	w_1	w_1	w_1	w_1	w_1	x_1	y_1	z_1	w_1

We note that (X, A) is a fuzzy lattice.

In Figure 2, we show the related tabular and graphical representation for the fuzzy relation A .

A	x_1	y_1	z_1	w_1
x_1	1.0	0.20	0.30	0.90
y_1	0.0	1.0	0.0	0.50
z_1	0.0	0.0	1.0	0.70
w_1	0.0	0.0	0.0	1.0




Figure 2

In Figure 3, we show the related tabular and graphical representations for the fuzzy relation A for $\alpha > 0.30$.

Here $x_1 \vee_\alpha w_1 = w_1$, $x_1 \wedge_\alpha w_1 = x_1$,

$y_1 \vee_\alpha w_1 = w_1$, $y_1 \wedge_\alpha w_1 = y_1$,

$z_1 \vee_\alpha w_1 = w_1$, $z_1 \wedge_\alpha w_1 = z_1$,

$y_1 \vee_\alpha z_1 = w_1$, $y_1 \vee_\alpha x_1 = w_1$, $z_1 \vee_\alpha x_1 = w_1$.

But $y_1 \wedge_\alpha z_1$, $y_1 \wedge_\alpha x_1$, $z_1 \wedge_\alpha x_1$ does not exist.




Figure 3

Remark 3.15. We note that Example 3.13 is an example of a fuzzy α -lattice for $\alpha = 0.2$ whereas Example 3.14, is not a fuzzy α -lattice for $\alpha > 0.30$.

Proposition 3.16. [4, Proposition 3.7] Let (X, A) be a fuzzy α -lattice, $\alpha \in [0, 1]$ and let $x, y, z \in X$. The following statements hold:

- (i) $A(x, x \vee_\alpha y) > \alpha$, $A(y, x \vee_\alpha y) > \alpha$, $A(x \wedge_\alpha y, x) > \alpha$, $A(x \wedge_\alpha y, y) > \alpha$;
- (ii) $A(x, z) > \alpha$ and $A(y, z) > \alpha$ implies $A(x \vee_\alpha y, z) > \alpha$;
- (iii) $A(z, x) > \alpha$ and $A(z, y) > \alpha$ implies $A(z, x \wedge_\alpha y) > \alpha$;
- (iv) $A(x, y) > \alpha$ iff $x \vee_\alpha y = y$;
- (v) $A(x, y) > \alpha$ iff $x \wedge_\alpha y = x$;
- (vi) If $A(y, z) > \alpha$, then $A(x \wedge_\alpha y, x \wedge_\alpha z) > \alpha$ and $A(x \vee_\alpha y, x \vee_\alpha z) > \alpha$;
- (vii) If $A(x \vee_\alpha y, z) > \alpha$, then $A(x, z) > \alpha$ and $A(y, z) > \alpha$;
- (viii) If $A(x, y \wedge_\alpha z) > \alpha$, then $A(x, y) > \alpha$ and $A(x, z) > \alpha$.

Proposition 3.17. [4, Proposition 3.8] Let (X, A) be a fuzzy α -lattice and let $x, y, z \in X$. Then

- (i) $x \vee_\alpha x = x$ and $x \wedge_\alpha x = x$;
- (ii) $x \vee_\alpha y = y \vee_\alpha x$ and $x \wedge_\alpha y = y \wedge_\alpha x$;
- (iii) $(x \vee_\alpha y) \vee_\alpha z = x \vee_\alpha (y \vee_\alpha z)$ and $(x \wedge_\alpha y) \wedge_\alpha z = x \wedge_\alpha (y \wedge_\alpha z)$;
- (iv) $(x \vee_\alpha y) \wedge_\alpha x = x$ and $(x \wedge_\alpha y) \vee_\alpha x = x$.

Lemma 3.18. [12, Lemma 3.18] Let (X, A) be a fuzzy α -lattice and $x, y, x', y' \in X$. If $A(x', x) > \alpha$ and $A(y', y) > \alpha$, then $A(x' \wedge_\alpha y', x \wedge_\alpha y) > \alpha$ and $A(x' \vee_\alpha y', x \vee_\alpha y) > \alpha$.

Definition 3.19. [4, Definition 3.8] Let (X, A) be a fuzzy α -lattice. (X, A) is fuzzy distributive iff $x \wedge_{\alpha} (y \vee_{\alpha} z) = (x \wedge_{\alpha} y) \vee_{\alpha} (x \wedge_{\alpha} z)$ and $(x \vee_{\alpha} y) \wedge_{\alpha} (x \vee_{\alpha} z) = x \vee_{\alpha} (y \wedge_{\alpha} z)$.

Note that (X, A) is fuzzy distributive iff $A(x \wedge_{\alpha} (y \vee_{\alpha} z), (x \wedge_{\alpha} y) \vee_{\alpha} (x \wedge_{\alpha} z)) > \alpha$ and $A((x \vee_{\alpha} y) \wedge_{\alpha} (x \vee_{\alpha} z), x \vee_{\alpha} (y \wedge_{\alpha} z)) > \alpha$.

Proposition 3.20. [12, Proposition 3.20] (*Modular inequality*) Let (X, A) be a fuzzy α -lattice and let $x, y, z \in X$. Then $A(x, z) > \alpha$ implies $A(x \vee_{\alpha} (y \wedge_{\alpha} z), (x \vee_{\alpha} y) \wedge_{\alpha} z) > \alpha$.

Definition 3.21. [12, Definition 3.21] Let (X, A) be a fuzzy α -lattice. (X, A) is fuzzy α -modular iff $A(x, z) > \alpha$ implies $x \vee_{\alpha} (y \wedge_{\alpha} z) = (x \vee_{\alpha} y) \wedge_{\alpha} z$ for all $x, y, z \in X$.

By the modular inequality, a fuzzy α -lattice (X, A) is fuzzy α -modular iff $A(x, z) > \alpha$ implies $A((x \vee_{\alpha} y) \wedge_{\alpha} z, x \vee_{\alpha} (y \wedge_{\alpha} z)) > \alpha$ for $x, y, z \in X$.

Proposition 3.22. [12, Proposition 3.22] Let (X, A) be a fuzzy α -lattice. (X, A) be a fuzzy distributive lattice, then (X, A) is fuzzy α -modular lattice.

We recall the Definition of a fuzzy α -modular pair in fuzzy α -lattice from paper [12]

Definition 3.23. [12, Definition 4.2] Let (X, A) be a fuzzy α -lattice. We say that (x, y) is a fuzzy α -modular pair and we write $(x, y)FM_{\alpha}$, if whenever $A(z, y) > \alpha$ for some $z \in X$, $\alpha \in [0, 1]$, then $(z \vee_{\alpha} x) \wedge_{\alpha} y = z \vee_{\alpha} (x \wedge_{\alpha} y)$.

We say that (x, y) is a fuzzy dual α -modular pair and we write $(x, y)FM_{\alpha}^*$, if whenever $A(y, z) > \alpha$ for some $z \in X$, then $(z \wedge_{\alpha} x) \vee_{\alpha} y = z \wedge_{\alpha} (x \vee_{\alpha} y)$.

We write $(x, y)\overline{FM}_{\alpha}$ when the pair (x, y) is not a fuzzy α -modular pair.

4. FUZZY α -MODULARITY IN FUZZY α -LATTICE

The following lemma gives a sufficient condition for a pair to be fuzzy α -modular in fuzzy α -lattice.

Lemma 4.1. If $A(x, y) > \alpha$ or $A(y, x) > \alpha$, then $(x, y)FM_{\alpha}$.

Proof. (i): Suppose that $A(x, y) > \alpha$ and $A(z, y) > \alpha$.

Then by Proposition 3.16(ii), we get

$$A(z \vee_{\alpha} x, y) > \alpha.$$

As $A(x, y) > \alpha$ by Proposition 3.16(v), we get

$$(4.1) \quad x \wedge_\alpha y = x.$$

We note that

$$\begin{aligned} & A((z \vee_\alpha x) \wedge_\alpha y, z \vee_F (x \wedge_\alpha y)) \\ &= A((z \vee_\alpha x) \wedge_\alpha y, z \vee_\alpha x), \quad \text{by (4.1)} \\ &= A(z \vee_\alpha x, z \vee_\alpha x), \quad \text{since } A(z \vee_\alpha x, y) > \alpha \\ &= 1 > 0 \end{aligned}$$

Therefore,

$$A((z \vee_\alpha x) \wedge_\alpha y, z \vee_F (x \wedge_\alpha y)) > \alpha.$$

We know that

$$A(z \vee_\alpha (x \wedge_\alpha y), (z \vee_\alpha x) \wedge_\alpha y) > \alpha$$

always holds.

By fuzzy antisymmetry of A we get

$$(z \vee_\alpha x) \wedge_\alpha y = z \vee_\alpha (x \wedge_\alpha y).$$

(ii): Suppose that $A(y, x) > \alpha$ and $A(z, y) > \alpha$.

By fuzzy transitivity of A we have

$$A(z, x) > \alpha.$$

We have

$$\begin{aligned} & A((z \vee_\alpha x) \wedge_\alpha y, z \vee_\alpha (x \wedge_\alpha y)) \\ & \geq \sup_{k \in X} \min[A((z \vee_\alpha x) \wedge_\alpha y, k), A(k, z \vee_F (x \wedge_\alpha y))] \\ & \geq \min[A((z \vee_\alpha x) \wedge_\alpha y, y), A(y, z \vee_\alpha (x \wedge_\alpha y))] \\ & \geq \min[A(x \wedge_\alpha y, y), A(y, z \vee_\alpha y)] \\ & \geq \min[A(y, y), A(y, y)] \end{aligned}$$

Therefore,

$$A((z \vee_\alpha x) \wedge_\alpha y, z \vee_\alpha (x \wedge_\alpha y)) > \alpha.$$

We know that

$$A(z \vee_\alpha (x \wedge_\alpha y), (z \vee_\alpha x) \wedge_\alpha y) > \alpha$$

always holds.

By fuzzy antisymmetry of A we get

$$(z \vee_\alpha x) \wedge_\alpha y = z \vee_\alpha (x \wedge_\alpha y).$$

Thus, $(x, y)FM_\alpha$ holds in either case. \square

Remark 4.2. If X has the elements \perp and \top , then for every $x \in X$, $(\perp, x)FM_\alpha$, $(x, \top)FM_\alpha$ and $(\perp, \top)FM_\alpha$ hold.

Remark 4.3. If $x, y \in X$, then, $(x \wedge_\alpha y, x)FM_\alpha$, $(x \wedge_\alpha y, y)FM_\alpha$, $(x, x \vee_\alpha y)FM_\alpha$, $(y, x \vee_\alpha y)FM_\alpha$ and $(x \wedge_\alpha y, x \vee_\alpha y)FM_\alpha$ hold.

We now prove some properties of fuzzy α -modular pairs.

Lemma 4.4. *If $(x, y)FM_\alpha$, $A(x \wedge_\alpha y, z) > \alpha$, then $(x \wedge_\alpha z, y)FM_\alpha$.*

Proof. Let $A(u, y) > \alpha$.

To show that $A([u \vee_\alpha (x \wedge_\alpha z)] \wedge_\alpha y, u \vee_\alpha [(x \wedge_\alpha z) \wedge_\alpha y]) > \alpha$ holds.
We know that

$$A(x \wedge_\alpha z, x) > \alpha.$$

By applying Proposition 3.16(vi), repeatedly we have

$$A(u \vee_\alpha (x \wedge_\alpha z), u \vee_\alpha x) > \alpha$$

and

$$(4.2) \quad A([u \vee_\alpha (x \wedge_\alpha z)] \wedge_\alpha y, (u \vee_\alpha x) \wedge_\alpha y) > \alpha.$$

As $(x, y)FM_\alpha$ holds so we have

$$(u \vee_\alpha x) \wedge_\alpha y = u \vee_\alpha (x \wedge_\alpha y).$$

Therefore, (4.2) reduces to

$$(4.3) \quad A([u \vee_\alpha (x \wedge_\alpha z)] \wedge_\alpha y, u \vee_\alpha (x \wedge_\alpha y)) > \alpha.$$

As $A(x \wedge_\alpha y, z) > \alpha$ we have

$$(x \wedge_\alpha y) \wedge_\alpha z = x \wedge_\alpha y.$$

Therefore, (4.3) reduces to

$$A([u \vee_\alpha (x \wedge_\alpha z)] \wedge_\alpha y, u \vee_\alpha [(x \wedge_\alpha y) \wedge_\alpha z]) > \alpha.$$

Thus,

$$A([u \vee_F (x \wedge_\alpha z)] \wedge_\alpha y, u \vee_\alpha [(x \wedge_\alpha z) \wedge_\alpha y]) > \alpha.$$

We know that

$$A(u \vee_\alpha [(x \wedge_\alpha z) \wedge_\alpha y], [u \vee_\alpha (x \wedge_\alpha z)] \wedge_\alpha y) > \alpha$$

always holds.

By fuzzy antisymmetry of A we get

$$[u \vee_\alpha (x \wedge_\alpha z)] \wedge_\alpha y = u \vee_\alpha [(x \wedge_\alpha z) \wedge_\alpha y].$$

Thus, $(x \wedge_\alpha z, y)FM_m$ holds. \square

Definition 4.5. Let $x, y \in X$. We say that (x, y) is a fuzzy independent pair and we write $(x, y) \perp FM_\alpha$ if $(x, y)FM_\alpha$ and $x \wedge_\alpha y = \perp$ hold.

Corollary 4.6. Let $x_1 \in X$. If $(x, y) \perp FM_\alpha$ and $A(x_1, x) > \alpha$, then $(x_1, y)FM_\alpha$.

Proof. Suppose that $(x, y) \perp FM_\alpha$ holds.

Then $(x, y)FM_\alpha$ holds with $x \wedge_\alpha y = \perp$.

As $A(\perp, x_1) > \alpha$ always holds we have

$$A(x \wedge_\alpha y, x_1) > \alpha.$$

Hence by Lemma 4.4, we have

$$(x \wedge_\alpha x_1, y)FM_\alpha.$$

As $A(x_1, x) > \alpha$, Proposition 3.16(v), we have

$$x \wedge_\alpha x_1 = x_1.$$

Thus, $(x_1, y)FM_\alpha$ holds. \square

Theorem 4.7. If $(x, y) \perp FM_\alpha$, $A(x_1, x) > \alpha$ and $A(y_1, y) > \alpha$, then $(x_1, y_1) \perp FM_\alpha$.

Proof. Suppose that $(x, y) \perp FM_\alpha$ holds.

Then $(x, y)FM_\alpha$ holds with $x \wedge_\alpha y = \perp$.

Let $A(x_1, x) > \alpha$ and $A(y_1, y) > \alpha$ for some $x_1, y_1 \in X$.

Then by Proposition 3.16(vi), we have

$$A(x_1 \wedge_\alpha y, x \wedge_\alpha y) > \alpha.$$

Therefore,

$$(4.4) \quad A(x_1 \wedge_\alpha y, \perp) > \alpha.$$

Similarly, $A(y_1, y) > \alpha$ by Proposition 3.16(vi), we have

$$(4.5) \quad A(x_1 \wedge_\alpha y_1, x_1 \wedge_\alpha y) > \alpha.$$

By fuzzy transitivity of A from (4.4) and (4.5) we get

$$(4.6) \quad A(x_1 \wedge_\alpha y_1, \perp) > \alpha.$$

As

$$(4.7) \quad A(\perp, x_1 \wedge_\alpha y_1) > \alpha$$

always holds.

From (4.6) and (4.7) by fuzzy antisymmetry of A we have

$$x_1 \wedge_\alpha y_1 = x_1 \wedge_\alpha y = \perp.$$

Now, it remains to show that $(x_1, y_1)FM_\alpha$ holds.

By Corollary 4.6, we have

$$(x_1, y)FM_\alpha.$$

Now, let $A(y_2, y_1) > \alpha$ for some $y_2 \in X$.

Then by (iv) and (v) of Proposition 3.16, we have

$$y_2 \vee_\alpha y_1 = y_1 \text{ and } y_2 \wedge_\alpha y_1 = y_2.$$

Since $A(y_2, y_1) > \alpha$ and $A(y_1, y) > \alpha$ by fuzzy transitivity of A we get

$$A(y_2, y) > \alpha.$$

As $A(y_1, y) > \alpha$, by (iv) and (v) of Proposition 3.16, we have

$$y_1 \vee_\alpha y = y \text{ and } y_1 \wedge_\alpha y = y_1.$$

Hence

$$\begin{aligned} & A((y_2 \vee_\alpha x_1) \wedge_\alpha y_1, y_2 \vee_\alpha (x_1 \wedge_\alpha y_1)) \\ &= A((y_2 \vee_\alpha x_1) \wedge_\alpha (y \wedge_\alpha y_1), y_2 \vee_\alpha (x_1 \wedge_\alpha y_1)) \\ &= A([(y_2 \vee_\alpha x_1) \wedge_\alpha y] \wedge_\alpha y_1, y_2 \vee_\alpha (x_1 \wedge_\alpha y_1)) \\ &= A([(y_2 \vee_\alpha (x_1 \wedge_\alpha y)] \wedge_\alpha y_1, y_2 \vee_\alpha (x_1 \wedge_\alpha y_1)), \text{ by } (x_1, y)FM_\alpha \\ &= A((y_2 \vee_\alpha \perp) \wedge_\alpha y_1, y_2 \vee_\alpha \perp) \\ &= A(y_2 \wedge_\alpha y_1, y_2) \\ &= A(y_2, y_2) = 1 > 0. \end{aligned}$$

Therefore,

$$A((y_2 \vee_\alpha x_1) \wedge_\alpha y_1, y_2 \vee_\alpha (x_1 \wedge_\alpha y_1)) > \alpha.$$

We know that

$$A(y_2 \vee_\alpha (x_1 \wedge_\alpha y_1), (y_2 \vee_\alpha x_1) \wedge_\alpha y_1) > \alpha$$

always holds.

By fuzzy antisymmetry of A we get

$$(y_2 \vee_\alpha x_1) \wedge_\alpha y_1 = y_2 \vee_\alpha (x_1 \wedge_\alpha y_1).$$

Thus, $(x_1, y_1)FM_\alpha$ holds.

Also, we have

$$x_1 \wedge_\alpha y_1 = \perp.$$

Hence $(x_1, y_1) \perp FM_\alpha$ holds. \square

Lemma 4.8. If $(x, y)FM_\alpha$ and if $(z, x \vee_\alpha y)FM_\alpha$, $A(z \wedge_\alpha (x \vee_\alpha y), x) > \alpha$, then $(z \vee_\alpha x, y)FM_\alpha$ and $(z \vee_\alpha x) \wedge_\alpha y = x \wedge_\alpha y$.

Proof. We have

$$\begin{aligned} & (z \vee_\alpha x) \wedge_\alpha y \\ &= (z \vee_\alpha x) \wedge_\alpha (x \vee_\alpha y) \wedge_\alpha y, \text{ by absorption identity} \\ &= (x \vee_\alpha z) \wedge_\alpha (x \vee_\alpha y) \wedge_\alpha y, \\ &= [x \vee_\alpha [z \wedge_\alpha (x \vee_\alpha y)]] \wedge_\alpha y, \text{ as } (z, x \vee_\alpha y)FM_\alpha \\ &= x \wedge_\alpha y, \text{ as } A(z \wedge_\alpha (x \vee_\alpha y), x) > 0. \end{aligned}$$

Thus, we get

$$(z \vee_\alpha x) \wedge_\alpha y = x \wedge_\alpha y.$$

We now show that $(z \vee_\alpha x, y)FM_\alpha$ holds, that is, to show that $A([y_1 \vee_\alpha (z \vee_\alpha x)] \wedge_\alpha y, y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]) > \alpha$. Let $A(y_1, y) > \alpha$ for some $y_1 \in X$.

We have

$$\begin{aligned} & A([y_1 \vee_\alpha (z \vee_\alpha x)] \wedge_\alpha y, y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]) \\ &= A([(y_1 \vee_\alpha x) \vee_\alpha z] \wedge_\alpha y, y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]) \\ &= A([(y_1 \vee_\alpha x) \vee_\alpha z] \wedge_\alpha (x \vee_\alpha y) \wedge_\alpha y, y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]), \\ & \quad \text{as } y = (x \vee_\alpha y) \wedge_\alpha y \\ &= A(y_1 \vee_\alpha [x \vee_F [z \wedge_\alpha (x \vee_\alpha y)]] \wedge_\alpha y, y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]) \\ &= A((y_1 \vee_\alpha x) \wedge_\alpha y, y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]), \\ & \quad \text{as } A(z \wedge_\alpha (x \vee_\alpha y), x) > \alpha \\ &= A(y_1 \vee_\alpha (x \wedge_\alpha y), y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]), \\ & \quad \text{as } (x, y)FM_m \\ &= A(y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y], y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]), \\ & \quad \text{as } x \wedge_\alpha y = (z \vee_\alpha x) \wedge_\alpha y \\ &= 1 > 0. \end{aligned}$$

Hence

$$A([y_1 \vee_\alpha (z \vee_\alpha x)] \wedge_\alpha y, y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]) > \alpha.$$

We know that

$$A(y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y], [y_1 \vee_\alpha (z \vee_\alpha x)] \wedge_\alpha y) > \alpha$$

always holds.

By fuzzy antisymmetry of A we get

$$[y_1 \vee_\alpha (z \vee_\alpha x)] \wedge_\alpha y = y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y].$$

Thus, $(z \vee_\alpha x, y)FM_\alpha$ holds. \square

Theorem 4.9. If $(x, y)FM_\alpha$ and $(z, x \vee_\alpha y) \perp FM_\alpha$, then $(z \vee_\alpha x, y)FM_\alpha$ and $(z \vee_\alpha x) \wedge_\alpha y = x \wedge_\alpha y$.

Proof. Suppose that $(x, y)FM_\alpha$ and $(z, x \vee_\alpha y) \perp FM_\alpha$ hold.

Then $(z, x \vee_\alpha y)FM_\alpha$ holds with $z \wedge_\alpha (x \vee_\alpha y) = \perp$.

Therefore, by Lemma 4.8, we have $(z \vee_\alpha x, y)FM_\alpha$ and $(z \vee_\alpha x) \wedge_\alpha y = x \wedge_\alpha y$. \square

Theorem 4.10. If $(x, y)FM_\alpha$ and $A(z, y) > \alpha$, then $(z \vee_\alpha x, y)FM_\alpha$.

Proof. Let $A(y_1, y) > \alpha$.

As $A(y_1, y) > \alpha$ and $A(z, y) > \alpha$ by Proposition 3.16(ii), we have

$$A(y_1 \vee_\alpha z, y) > \alpha.$$

Also, $(x, y)FM_\alpha$ holds so we have

$$(4.8) \quad [(y_1 \vee_\alpha z) \vee_\alpha x] \wedge_\alpha y = (y_1 \vee_\alpha z) \vee_\alpha (x \wedge_\alpha y).$$

To show that $A([y_1 \vee_\alpha (z \vee_\alpha x)] \wedge_\alpha y, y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_F y]) > \alpha$.

Consider

$$\begin{aligned} & A([y_1 \vee_\alpha (z \vee_\alpha x)] \wedge_\alpha y, y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]) \\ &= A([(y_1 \vee_\alpha z) \vee_\alpha x] \wedge_\alpha y, y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]) \\ &= A((y_1 \vee_\alpha z) \vee_\alpha (x \wedge_\alpha y), y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]), \quad \text{by (4.8)} \\ &= A(y_1 \vee_\alpha [z \vee_\alpha (x \wedge_\alpha y)], y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]) \\ &= A(y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y], y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]), \quad \text{as } (x, y)FM_\alpha \\ &= 1 > 0 \end{aligned}$$

Hence

$$A([y_1 \vee_\alpha (z \vee_\alpha x)] \wedge_\alpha y, y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y]) > \alpha.$$

We know that

$$A(y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y], [y_1 \vee_\alpha (z \vee_\alpha x)] \wedge_\alpha y) > \alpha$$

always holds.

By fuzzy antisymmetry of A we get

$$[y_1 \vee_\alpha (z \vee_\alpha x)] \wedge_\alpha y = y_1 \vee_\alpha [(z \vee_\alpha x) \wedge_\alpha y].$$

Thus, $(z \vee_{\alpha} x, y)FM_{\alpha}$ holds. \square

Corollary 4.11. *If $(x, y) \perp FM_{\alpha}$ and $A(z, y) > \alpha$, then $(z \vee_{\alpha} x, y)FM_{\alpha}$ and $(z \vee_{\alpha} x) \wedge_{\alpha} y = z$.*

Proof. Suppose that $(x, y) \perp FM_{\alpha}$ holds.

Then $(x, y)FM_{\alpha}$ holds with $x \wedge_{\alpha} y = \perp$.

Also, given $A(z, y) > \alpha$.

Therefore, by Theorem 4.10, we have

$$(z \vee_{\alpha} x, y)FM_{\alpha}.$$

Now, it remains to show that $(z \vee_{\alpha} x) \wedge_{\alpha} y = z$.

By $(x, y)FM_{\alpha}$ and $A(z, y) > \alpha$ we have

$$(z \vee_{\alpha} x) \wedge_{\alpha} y = z \vee_{\alpha} (x \wedge_{\alpha} y) = z \vee_{\alpha} \perp = z. \quad \square$$

Lemma 4.12. *If $(x, y) \perp FM_{\alpha}$ and $(z, x \vee_{\alpha} y) \perp FM_{\alpha}$, then $(z \vee_{\alpha} x, y) \perp FM_{\alpha}$.*

Proof. Suppose that $(x, y) \perp FM_{\alpha}$ and $(z, x \vee_{\alpha} y) \perp FM_{\alpha}$ hold.

Then $(x, y)FM_{\alpha}$ and $(z, x \vee_{\alpha} y)FM_{\alpha}$ hold with

$$x \wedge_{\alpha} y = \perp \text{ and } z \wedge_{\alpha} (x \vee_{\alpha} y) = \perp.$$

By Theorem 4.9, we get

$$(z \vee_{\alpha} x, y)FM_{\alpha} \text{ and } (z \vee_{\alpha} x) \wedge_{\alpha} y = x \wedge_{\alpha} y = \perp.$$

Hence $(z \vee_{\alpha} x, y) \perp FM_{\alpha}$ holds. \square

Definition 4.13. Let $\mathcal{L} = (X, A)$ be a fuzzy α -lattice. Let $x, y \in X$, then $y \prec_F^{\alpha} x$ (x “fuzzy covers” y) if $\alpha < A(y, x) < 1$, $A(y, c) > \alpha$ and $A(c, x) > \alpha$ imply $c = y$ or $c = x$.

Definition 4.14. Let P denote the set of all $x \in X$ such that $\perp \prec_F^{\alpha} x$. The elements of P are called fuzzy atoms.

Corollary 4.15. *Let $\mathcal{L} = (X, A)$ be a fuzzy α -lattice with \perp . If $p \in P$, $y \in X$, then $(y, p)FM_{\alpha}$.*

Proof. If $A(x, p) > \alpha$, then $x = \perp$ or $x = p$.

Case (1): If $x = \perp$, then

$$(x \vee_{\alpha} y) \wedge_{\alpha} p = (\perp \vee_{\alpha} y) \wedge_{\alpha} p = y \wedge_{\alpha} p = x \vee_{\alpha} (y \wedge_{\alpha} p).$$

Case (2): If $x = p$, then

$$(x \vee_{\alpha} y) \wedge_{\alpha} p = (p \vee_{\alpha} y) \wedge_{\alpha} p = p = p \vee_{\alpha} (y \wedge_{\alpha} p) = x \vee_{\alpha} (y \wedge_{\alpha} p).$$

Thus, $(y, p)FM_{\alpha}$ holds. \square

5. FUZZY SEMI-MODULAR IN α -LATTICES

In this section, we introduce the notion of a fuzzy semi-modular fuzzy α -lattice.

Definition 5.1. A fuzzy α -lattice $\mathcal{L} = (X, A)$ with \perp is called fuzzy weakly α -modular when in $\mathcal{L} = (X, A)$, $x \wedge_\alpha y \neq \perp$ implies $(x, y)FM_\alpha$.

Definition 5.2. A fuzzy α -lattice (X, A) with \perp is called \perp_F -symmetric fuzzy α -lattice when in (X, A) , $(x, y) \perp FM_\alpha$ implies $(y, x)FM_\alpha$.

Definition 5.3. A fuzzy weakly modular α -lattice with \perp_F -symmetric fuzzy α -lattice is called as a fuzzy semi-modular α -lattice.

Throughout this section, we assume $\mathcal{L} = (X, A)$ as a fuzzy semi-modular α -lattice.

Lemma 5.4. If $x \wedge_\alpha y \prec_F^\alpha x$, then $y \prec_F^\alpha x \vee_\alpha y$.

Proof. Suppose that $A(y, z) > \alpha$ and

$$(5.1) \quad A(z, x \vee_\alpha y) > \alpha.$$

To show that $y = z$ or $x \vee_F y = z$.

Define $u = z \wedge_\alpha x$.

Then

$$A(x \wedge_\alpha y, u) > \alpha \text{ and } A(u, x) > \alpha.$$

Hence

$$x \wedge_\alpha y = u \text{ or } u = x \text{ as } x \wedge_\alpha y \prec_F^\alpha x.$$

Case (1): If $u = x$, then $z \wedge_\alpha x = x$,
that is, $A(x, z) > \alpha$ by Proposition 3.16(v).
So, by Proposition 3.16(vi), we have

$$A(x \vee_\alpha y, z \vee_\alpha y) > \alpha.$$

Therefore, by (5.1) we get

$$(5.2) \quad A(x \vee_\alpha y, z) > \alpha.$$

From (5.1) and (5.2), by fuzzy antisymmetry of A we get

$$x \vee_\alpha y = z.$$

Case (2): Let $u = x \wedge_\alpha y$, i.e., $z \wedge_\alpha x = x \wedge_\alpha y$.

Now, if $x \wedge_\alpha y \neq \perp$, then $z \wedge_\alpha x \neq \perp$.

By the definition of fuzzy weakly modular α -lattice we have $(x, z)FM_\alpha$.

If $x \wedge_{\alpha} z = x \wedge_{\alpha} y = \perp$, then $\perp \prec_F^{\alpha} x$,
that is, $x \in P$ and $(z, x)FM_{\alpha}$ by Corollary 4.15.

Thus we have $(x, z)FM_{\alpha}$ as \mathcal{L} is \perp_F -symmetric fuzzy α -lattice.
Now, $(x, z)FM_{\alpha}$ and $A(y, z) > \alpha$ imply that

$$z = (y \vee_{\alpha} x) \wedge_{\alpha} z = y \vee_{\alpha} (x \wedge_{\alpha} z) = y \vee_{\alpha} (x \wedge_{\alpha} y) = y \vee_{\alpha} \perp = y.$$

From Case (1) and Case (2) we have either

$$y = z \text{ or } z = x \vee_{\alpha} y.$$

Therefore, $y \prec_F^{\alpha} x \vee_{\alpha} y$. \square

Lemma 5.5. *If $y \prec_F^{\alpha} x \vee_{\alpha} y$ and if $(y, x)FM_{\alpha}$, then $x \wedge_{\alpha} y \prec_F^{\alpha} x$.*

Proof. If $x \wedge_{\alpha} y = x$, then $x \vee_{\alpha} y = y$, contrary to $y \prec_F^{\alpha} x \vee_{\alpha} y$.

Hence $\alpha < A(x \wedge_{\alpha} y, x) < 1$.

Now, suppose that

$$A(x \wedge_{\alpha} y, z) > \alpha$$

and

$$(5.3) \quad A(z, x) > \alpha.$$

Define $u = z \vee_{\alpha} y$.

Then $A(y, u) > \alpha$ and $A(u, x \vee_{\alpha} y) > \alpha$.

Hence $u = y$ or $u = x \vee_{\alpha} y$ as $y \prec_F^{\alpha} x \vee_{\alpha} y$.

Case (1): If $u = y$, then $y = z \vee_{\alpha} y$,

that is, $A(z, y) > \alpha$ by Proposition 3.16(iv).

Therefore, by Proposition 3.16(vi), we get

$$(5.4) \quad A(z \wedge_{\alpha} x, y \wedge_{\alpha} x) > \alpha.$$

As $A(z, x) > \alpha$ so by Proposition 3.16(v), we have

$$z \wedge_{\alpha} x = z.$$

Therefore, (5.4) reduces to

$$(5.5) \quad A(z, y \wedge_{\alpha} x) > \alpha.$$

Hence from (5.3) and (5.5) by fuzzy antisymmetry of A we get

$$x \wedge_{\alpha} y = z.$$

Case (2): On the other hand if $u = x \vee_{\alpha} y$, then $z \vee_{\alpha} y = x \vee_{\alpha} y$.

Hence by $(y, x)FM_{\alpha}$ we get

$$x = (x \vee_{\alpha} y) \wedge_{\alpha} x = (z \vee_{\alpha} y) \wedge_{\alpha} x = z \vee_{\alpha} (y \wedge_{\alpha} x) = z.$$

Hence from Case (1) and Case (2) we have either

$$x \wedge_{\alpha} y = z \text{ or } z = x.$$

Thus, $x \wedge_{\alpha} y \prec_F^{\alpha} x$ holds. \square

Lemma 5.6. *If $x \prec_F^{\alpha} y$ and $z \in X$, then either*

- (i) $x \vee_{\alpha} z = y \vee_{\alpha} z$ or
- (ii) $x \vee_{\alpha} z \prec_F^{\alpha} y \vee_{\alpha} z$.

Proof. Clearly $x \vee_{\alpha} z = y \vee_{\alpha} z$ or $\alpha < A(x \vee_{\alpha} z, y \vee_{\alpha} z) < 1$.

Suppose that $A(x \vee_{\alpha} z, u) > \alpha$ and $A(u, y \vee_{\alpha} z) > \alpha$.

Then by Proposition 3.16(vi), we get

$$A((x \vee_{\alpha} z) \wedge_{\alpha} y, u \wedge_{\alpha} y) > \alpha \text{ and } A(u \wedge_{\alpha} y, (y \vee_{\alpha} z) \wedge_{\alpha} y) > \alpha,$$

i.e.,

$$A((x \vee_{\alpha} z) \wedge_{\alpha} y, u \wedge_{\alpha} y) > \alpha, \text{ and } A(u \wedge_{\alpha} y, y) > \alpha.$$

As $A(x, x \vee_{\alpha} z) > \alpha$ always holds by Proposition 3.16(vi), we get

$$A(x \wedge_{\alpha} y, (x \vee_{\alpha} z) \wedge_{\alpha} y) > \alpha.$$

As $x \prec_F^{\alpha} y$ we get

$$(5.6) \quad A(x, (x \vee_{\alpha} z) \wedge_{\alpha} y) > \alpha.$$

Also,

$$(5.7) \quad A((x \vee_{\alpha} z) \wedge_{\alpha} y, y \wedge_{\alpha} u) > \alpha.$$

From (5.6) and (5.7) by fuzzy transitivity of A we get

$$A(x, y \wedge_{\alpha} u) > \alpha$$

and

$$A(y \wedge_{\alpha} u, y) > \alpha$$

always holds.

If $y \wedge_{\alpha} u = \perp$, then for $x = \perp$ and $y \in P$ we get

$$(y, u)FM_{\alpha}.$$

If $y \wedge_{\alpha} u \neq \perp$, then $(y, u)FM_{\alpha}$ by the definition of fuzzy weakly α -modular.

Therefore, we get $(y, u)FM_{\alpha}$ in either case.

Hence

$$z \vee_{\alpha} (y \wedge_{\alpha} u) = (z \vee_{\alpha} y) \wedge_{\alpha} u = u.$$

Since $A(z, x \vee_{\alpha} z) > \alpha$ and $A(x \vee_{\alpha} z, u) > \alpha$.

Now, since $x \prec_F^{\alpha} y$, we have

$$x = y \wedge_{\alpha} u \text{ or } y \wedge_{\alpha} u = y.$$

If $y \wedge_{\alpha} u = x$, then

$$z \vee_{\alpha} x = z \vee_{\alpha} (y \wedge_{\alpha} u) = u,$$

if $y \wedge_{\alpha} u = y$, then

$$z \vee_{\alpha} y = z \vee_{\alpha} (y \wedge_{\alpha} u) = u.$$

This shows that either

$$x \vee_{\alpha} z = u \text{ or } u = y \vee_{\alpha} z.$$

Hence we have

$$x \vee_{\alpha} z \prec_F^{\alpha} y \vee_{\alpha} z.$$

Thus, (ii) holds. \square

Lemma 5.7. *If $y \prec_F^{\alpha} z$, $(x, z)FM_{\alpha}$ and $(x, y)FM_{\alpha}$, then either*

- (i) $x \vee_{\alpha} y \prec_F^{\alpha} x \vee_{\alpha} z$ and $x \wedge_{\alpha} y = x \wedge_{\alpha} z$ or
- (ii) $x \vee_{\alpha} y = x \vee_{\alpha} z$ and $x \wedge_{\alpha} y \prec_F^{\alpha} x \wedge_{\alpha} z$.

Proof. As $(x, z)FM_{\alpha}$ holds, we have

$$(y \vee_{\alpha} x) \wedge_{\alpha} z = y \vee_{\alpha} (x \wedge_{\alpha} z).$$

Let $u = (y \vee_{\alpha} x) \wedge_{\alpha} z = y \vee_{\alpha} (x \wedge_{\alpha} z)$.

Then by (iv) and (v) of Proposition 3.16 we have

$$A(y, u) > \alpha \text{ and } A(u, z) > \alpha.$$

As $y \prec_F^{\alpha} z$ either $y = u$ or $u = z$.

Case (1): Suppose that $y = u$.

Then $y = y \vee_{\alpha} (x \wedge_{\alpha} z)$, by Proposition 3.16(iv) we get

$$A(x \wedge_{\alpha} z, y) > \alpha.$$

By Proposition 3.16(vi), we get

$$(5.8) \quad A(x \wedge_{\alpha} z, y \wedge_{\alpha} x) > \alpha.$$

As $y \prec_F^{\alpha} z$ we have $\alpha < A(y, z) < 1$. Hence

$$(5.9) \quad A(x \wedge_{\alpha} y, x \wedge_{\alpha} z) > \alpha.$$

From (5.8) and (5.9) by fuzzy antisymmetry of A we get

$$x \wedge_{\alpha} z = x \wedge_{\alpha} y.$$

Moreover $u \prec_F^{\alpha} z$, that is,

$$(x \vee_{\alpha} y) \wedge_{\alpha} z \prec_F^{\alpha} z.$$

Hence by Lemma 5.4, we get

$$(5.10) \quad x \vee_{\alpha} y \prec_F^{\alpha} (x \vee_{\alpha} y) \vee_{\alpha} z.$$

As $\alpha < A(y, z) < 1$ by Proposition 3.16(vi), we get $y \vee_{\alpha} z = z$. Therefore (5.10) reduces to

$$x \vee_{\alpha} y \prec_F^{\alpha} x \vee_{\alpha} z.$$

Thus, (i) holds.

Case (2): Now let us suppose that $u = z$.

Then $(y \vee_{\alpha} x) \wedge_{\alpha} z = z$, by Proposition 3.16(iv) we get

$$A(z, y \vee_{\alpha} x) > \alpha.$$

By Proposition 3.16(vi), we get

$$A(x \vee_{\alpha} z, y \vee_{\alpha} x) > \alpha.$$

Also, $\alpha < A(y, z) < 1$ by Proposition 3.16(vi), we have

$$A(x \vee_{\alpha} y, x \vee_{\alpha} z) > \alpha.$$

Thus, by fuzzy antisymmetry of A we get

$$x \vee_{\alpha} z = x \vee_{\alpha} y.$$

Now, $y \prec_F^{\alpha} z = u = y \vee_{\alpha} (x \wedge_{\alpha} z) = (y \vee_{\alpha} x) \wedge_{\alpha} z$.

Now, $\alpha < A(y, z) < 1$ by Proposition 3.16(vi), we have

$$A(x \wedge_{\alpha} y, x \wedge_{\alpha} z) > \alpha.$$

As $A(x \wedge_{\alpha} z, z) > \alpha$ always holds, so by fuzzy transitivity of A we have

$$A(x \wedge_{\alpha} y, z) > \alpha.$$

Since $(x, y)_F M_m$, $A(x \wedge_{\alpha} y, z) > \alpha$, then by Lemma 4.4, we have

$$(x \wedge_{\alpha} z, y)_F M_m.$$

Thus, by Lemma 5.5, we get

$$x \wedge_{\alpha} z \wedge_{\alpha} y \prec_F^{\alpha} x \wedge_{\alpha} z,$$

or equivalently,

$$x \wedge_{\alpha} y \prec_F^{\alpha} x \wedge_{\alpha} z.$$

Thus, (ii) holds. \square

6. Conclusion

In this paper, we have studied the notion of a fuzzy independent pair and obtained some properties of fuzzy α -modular pairs and independent pairs in fuzzy α -lattice.

7. Acknowledgement

The authors are thankful to the referee for fruitful suggestions, which enhanced the quality of the paper.

REFERENCES

- [1] N. Ajmal and K. V. Thomas, *Fuzzy lattices*, Information Science, **79** (1994), 271–291.
- [2] I. Chon, *Fuzzy partial order relations and fuzzy lattices*, Korean J. Math., **17(4)** (2009), 361–374.
- [3] F. Maeda and S. Maeda, *Theory of Symmetric Lattices*, Springer-verlag, Berlin, 1970.
- [4] I. Mezzomo and B. Bedregal, *On fuzzy α -lattices*, IEEE World Congress on Computational Intelligence - WCCI (2016), 775–781. doi:10.1109/FUZZ-IEEE.2016.7737766.
- [5] S. K. Nimborkar and J. A. Khubchandani, *Fuzzy essential-small sub-modules and fuzzy small-essential submodules*, Journal of Hyperstructures, **9(2)** (2020), 52–67.
- [6] S. K. Nimborkar and J. A. Khubchandani, *Fuzzy weakly irreducible ideals of a ring*, Journal of Hyperstructures, **10(2)** (2021), 118–128.
- [7] M. P. Wasadikar and P. A. Khubchandani, *Fuzzy modularity in fuzzy lattices*, The Journal of Fuzzy Mathematics, **27(4)** (2019), 985–998.
- [8] M. P. Wasadikar and P. A. Khubchandani, *Fuzzy modularity and complement in fuzzy lattices*, TWMS J. App. and Eng. Math., **12(4)** (2022), 1368–1379.
- [9] M. P. Wasadikar and P. A. Khubchandani, *Fuzzy semi-orthogonality in fuzzy lattices*, (Communicated).
- [10] M. P. Wasadikar and P. A. Khubchandani, ‘*Del’ Relation and parallelism in fuzzy lattices*, TWMS J. App. and Eng. Math., **12(3)** (2022), 864–875.
- [11] M. P. Wasadikar and P. A. Khubchandani, *Fuzzy distributive pairs in fuzzy lattices* Discussions Mathematicae General Algebra and Applications, **42(1)** (2022), 179–199.
- [12] M. P. Wasadikar and P. A. Khubchandani, *Fuzzy pairs in fuzzy α -lattices*, Journal of Hyper structures, **9(2)** (2020), 68–80.
- [13] B. Yaun and W. Wu, *Fuzzy ideals on distributive lattices*, Fuzzy sets and Systems, **35** (1990), 231–240.
- [14] L. Zadeh, *Fuzzy sets*, Information and control, **8** (1965), 338–353.
- [15] L. Zadeh, *Similarity relations and fuzzy orderings*, Information Sciences, **3** (1971), 177–200.

Payal A. Khubchandani

Department of Engineering Sciences and Humanities,
Vishwakarma Institute of Technology, Pune 411037, India
Email: payal_khubchandani@yahoo.com

Jyoti A. Khubchandani

Department of Engineering Sciences and Humanities,
Vishwakarma Institute of Technology, Pune 411037, India
Email: khubchandani_jyoti@yahoo.com