



## Research Paper

## COMPUTATION OF NM-POLYNOMIAL AND TOPOLOGICAL INDICES FOR CYCLE RELATED GRAPHS

DEEPASREE SASI KUMAR<sup>1</sup> , RANJINI<sup>2</sup> , HARINI<sup>3</sup> VEERABHADRAIAH LOKESHA<sup>4,\*</sup> <sup>1</sup>Acharya Institute of Technology, Bangalore, affiliated to Visvesvaraya Technological University, Belagavi, Karnataka, India, drdeepasreeskumar@gmail.com<sup>2</sup>Department of Mathematics, Don Bosco Institute Of Technology, Kumbalagodu, Bangaluru-56, Karnataka, India, drranjinips@gmail.com<sup>3</sup>Department of Mathematics, PES College of Engineering, Mandya, Karnataka, India, harinimrmaha@gmail.com<sup>4</sup>DOS in Mathematics, Vijyanagara Sri Krishnadevaraya University, Ballari-583105, India, v.lokesha@gmail.com

## ARTICLE INFO

## Article history:

Received: 10 June 2025

Accepted: 17 August 2025

Communicated by Mahmood Bakhshi

## Keywords:

Topological indices

wheel graph

helm graph

gear graph

## MSC:

05C09, 05C07, 05C92.

## ABSTRACT

This paper focuses on the computation of  $NM$ -polynomial and several topological indices for cycle related graphs such as Wheel graph, Helm graph and Gear graph. The  $NM$ -polynomial is a graph invariant that encodes information about the sub graph structure, which is crucial for understanding the connectivity and combinatorial properties of a graph. We develop formulas and methods for computing the  $NM$ -polynomial for specific cycle-related graphs, demonstrating its utility in capturing key graph characteristics.

## 1. INTRODUCTION

Topological indices are numerical descriptors that describe a molecular graph's topology, or structure. In the study of quantitative structure-activity relationships (QSAR) and quantitative structure-property relationships (QSPR), these indices are particularly useful in chemistry [1, 3, 18, 19, 20, 22]. Topological indices are essential for comprehending the

\*Address correspondence to Deepasree Sasi Kumar; Department of Mathematics, Acharya Institute of Technology, Bangalore, affiliated to Visvesvaraya Technological University, Belagavi, Karnataka, India, E-mail: drdeepasreeskumar@gmail.com.

features and behaviors of cycle-related graphs, which comprise chemical structures with one or more rings. Among different types of indices, degree based indices are derived from the degrees of the vertices of a molecular graph [5, 8, 9, 10, 11, 12]. Firstly, Wiener index is introduced by the chemist Harold Wiener in 1947 [23] and is defined as the sum of the distances between all unordered pairs of vertices of a graph  $G$ , that is

$$W(G) = \sum_{u,v \in V(G)} d_G(u,v),$$

where  $V(G)$  is the vertex set of  $G$  and  $d_G(u,v)$  denotes distance from  $u$  to  $v$ .

In recent years, the application of topological indices has extended beyond traditional QSAR and QSPR studies into areas such as nanotechnology, pharmacology, and materials science. These indices serve as effective mathematical tools for predicting the stability, reactivity, and biological activity of chemical compounds without the need for exhaustive experimental procedures. Furthermore, with the advancement of computational techniques, more sophisticated and generalized forms of topological indices have been developed to capture subtle structural variations within molecular graphs. The study of cycle-containing graphs, such as benzenoid systems and fullerene structures, has especially benefited from these indices, enabling researchers to establish correlations between molecular topology and physicochemical properties with greater precision.

Researchers have recently focused on neighborhood degree-based indices [6, 13, 14, 21]. The relations of some neighborhood degree-based topological indices with the  $NM$ -polynomial [4, 7, 15, 17] are shown in Table 1

**Definition 1.1.** Let  $G$  be a graph. Then  $NM$ -Polynomial [13, 16] of  $G$  is defined as

$$(1.1) \quad NM(G : x, y) = \sum_{i \leq j} m_{ij}^*(G) x^i y^j.$$

where  $m_{ij}^*(G)$ ,  $i, j \geq 1$  be the number of edges  $uv$  of  $G$  such that  $\delta_u, \delta_v = \{i, j\}$ . Here  $\delta_u, \delta_v$  represent the neighborhood degree sum of the vertices  $u$  and  $v$  in  $G$ .

TABLE 1. Expression of TI's using  $NM$ -polynomial [2]

| TI's         | $f(s, t)$              | Expression with $NM$ -Polynomial          |
|--------------|------------------------|-------------------------------------------|
| $M_1'$       | $s + t$                | $(D_s + D_t)NM(G : s, t)_{s=t=1}$         |
| $M_2^*$      | $st$                   | $(D_s D_t)NM(G : s, t)_{s=t=1}$           |
| ${}^{nm}M_2$ | $\frac{1}{st}$         | $(S_s S_t)NM(G : s, t)_{s=t=1}$           |
| $NSDD$       | $\frac{s^2 + t^2}{st}$ | $(D_s S_t + D_t S_s)NM(G : s, t)_{s=t=1}$ |
| $NH$         | $\frac{2}{s+t}$        | $2S_s JNM(G; s, t)_{s=t=1}$               |
| $NI$         | $\frac{2}{s+t}$        | $S_s J D_s D_t NM(G; s, t)_{s=t=1}$       |

Where  $D_x = x \frac{\partial}{\partial x} [M(G)]$ ,  $D_y = y \frac{\partial}{\partial y} [M(G)]$ ,  $D_x D_y = x_1 \frac{\partial}{\partial x} (D_y)$ ,

$$S_x = \int_0^x \frac{M(G; t, y)}{t} dt, S_y = \int_0^y \frac{M(G; x, t)}{t} dt, S_x S_y = \int_0^x \frac{S_y(t, y)}{t} dt.$$

## 2. RESULTS AND DISCUSSION

In this section, we obtained  $NM$ -polynomial and some topological indices of cycle related graphs such as wheel graph, helm graph and gear graph.

**2.1. Wheel graph.** Wheel graph is a graph formed by connecting a single universal vertex to all vertices of a cycle.

Let  $W_n$  be the wheel graph with  $n + 1$  number of vertices and  $2n$  number of edges. The edge partition of neighborhood degree sequences of  $W_n$  is given by

$$(2.1) \quad E_{(6+n)(3n)} = n, E_{(6+n)(6+n)} = n$$

**Theorem 2.1.**  $NM$ -polynomial of wheel graph is given by,

$$NM(W_n : s, t) = ns^{n+6} [t^{3n} + t^{n+6}].$$

*Proof.* By using equation (2.1)

$$\begin{aligned} NM(W_n : s, t) &= m_{(n+6)3n} s^{n+6} t^{3n} + m_{(n+6)(n+6)} s^{n+6} t^{n+6} \\ &= n s^{n+6} t^{3n} + n s^{n+6} t^{n+6} \\ &= ns^{n+6} [t^{3n} + t^{n+6}]. \end{aligned}$$

□

**Theorem 2.2.** Let  $W_n$  be the wheel graph, then

- (1) Neighborhood First Zagreb index,  $M_1'(W_n : s, t) = 6n(n + 3)$ .
- (2) Neighborhood Second Zagreb index,  $M_2^*(W_n : s, t) = 2n(2n^2 + 15n + 18)$ .
- (3) Neighborhood Modified Second Zagreb index,  ${}^{nm}M_2(W_n : s, t) = \frac{2(2n + 3)}{(n + 6)^2}$ .
- (4) Neighborhood Symmetric division deg index,  $NSDD(W_n : s, t) = \frac{(4n + 6)^2}{3(n + 6)}$ .
- (5) Neighborhood Harmonic Index,  $NH(W_n : s, t) = \frac{3n(n + 3)}{(n + 6)(2n + 3)}$ .
- (6) Neighborhood Inverse Index,  $NI(W_n : s, t) = \frac{n(n + 6)(5n + 3)}{2(2n + 3)}$ .

*Proof.* Using equation (2.1) and Table 1

$$\begin{aligned} (1) D_s[NM(W_n)] &= s \frac{\partial}{\partial s} [NM(W_n)] = sn(n + 6)s^{n+5} [t^{3n} + t^{n+6}] \\ &= n(n + 6)s^{n+6} [t^{3n} + t^{n+6}] \end{aligned}$$

$$\begin{aligned} D_t[NM(W_n)] &= t \frac{\partial}{\partial t} [NM(W_n)] = tns^{n+6} [3nt^{3n-1} + (n + 6)t^{n+5}] \\ &= ns^{n+6} [3nt^{3n} + (n + 6)t^{n+6}] \end{aligned}$$

$$M_1'(W_n) = (D_s + D_t) [NM(W_n; s, t)]_{s=t=1}$$

$$(D_s + D_t)_{s=t=1} = 6(n^2 + 3n) = 6n(n + 3).$$

$$\begin{aligned} (2) \quad D_s D_t [NM(W_n)] &= s \frac{\partial}{\partial s} [D_t] = sn(n + 6)s^{n+5} [3nt^{3n} + (n + 6)t^{n+6}] \\ &= s [n(n + 6)s^{n+5} [3nt^{3n} + (n + 6)t^{n+6}]] \\ &= 3n^2(n + 6)s^{n+6} t^{3n} + n(n + 6)^2 s^{n+6} t^{n+6} \end{aligned}$$

$$\begin{aligned} M_2^*(W_n) &= (D_s D_t) [NM(W_n; s, t)]_{s=t=1} = 3n^2(n + 6) + n(n + 6)^2 \\ &= 2n(2n^2 + 15n + 18). \end{aligned}$$

$$\begin{aligned} (3) \quad S_s &= \int_0^s \frac{NM(W_n; x, t)}{x} dx \\ &= \int_0^s nx^{n+5} [t^{3n} + t^{n+6}] dx \\ &= n \frac{s^{n+6}}{n + 6} [t^{3n} + t^{n+6}] \end{aligned}$$

$$\begin{aligned} S_t &= \int_0^t \frac{NM(W_n; s, y)}{y} dy \\ &= \int_0^t ns^{n+6} [y^{3n-1} + y^{n+5}] dy \\ &= ns^{n+6} \left[ \frac{t^{3n}}{3n} + \frac{t^{n+6}}{n + 6} \right] \end{aligned}$$

$$\begin{aligned} S_s S_t &= \int_0^s \frac{S_t(x, t)}{x} dx \\ &= \int_0^s nx^{n+5} \left[ \frac{t^{3n}}{3n} + \frac{t^{n+6}}{n + 6} \right] dx \\ &= n \frac{s^{n+6}}{n + 6} \left[ \frac{t^{3n}}{3n} + \frac{t^{n+6}}{n + 6} \right] \end{aligned}$$

$$\begin{aligned} {}^{nm}M_2(W_n : s, t) &= (S_s S_t) [NM(W_n; s, t)]_{s=t=1} \\ &= \frac{2(2n + 3)}{(n + 6)^2}. \end{aligned}$$

$$(4) \quad NSDD(W_n : s, t) = [D_s S_t + S_s D_t] [NM(W_n; s, t)]_{s=t=1}$$

$$\begin{aligned} D_s S_t &= s \frac{\partial}{\partial s} (S_t) \\ &= s \frac{\partial}{\partial s} \left[ n s^{n+6} \left( \frac{t^{3n}}{3n} + \frac{t^{n+6}}{n+6} \right) \right] \\ &= n(n+6) s^{n+6} \left[ \frac{t^{3n}}{3n} + \frac{t^{n+6}}{n+6} \right] \end{aligned}$$

$$\begin{aligned} S_s D_t &= \int_0^s \frac{D_t(x, t)}{x} dx \\ &= \int_0^s n x^{n+5} [3n t^{3n} + (n+6) t^{n+6}] dx \\ &= n \frac{s^{n+6}}{n+6} [3n t^{3n} + (n+6) t^{n+6}] \end{aligned}$$

$$\begin{aligned} NSDD(W_n) &= n(n+6) \left[ \frac{1}{3n} + \frac{1}{n+6} \right] + \frac{n}{n+6} [3n + (n+6)] \\ &= \frac{(4n+6)^2}{3(n+6)}. \end{aligned}$$

$$(5) \quad NH(W_n : S, t) = 2S_s J [NM(W_n : s, t)]_{s=t=1}$$

$$\begin{aligned} J[NM(W_n; s, t)] &= NM(W_n; s, s) \\ &= n s^{n+6} [s^{3n} + s^{n+6}] \\ &= n [s^{4n+6} + s^{2n+12}] \end{aligned}$$

$$\begin{aligned} S_s J [NM(W_n)] &= \int_0^s \frac{J [NM(W_n)]_x}{x} dx \\ &= n \int_0^s (x^{4n+5} + x^{2n+11}) dx \\ &= n \left[ \frac{s^{4n+6}}{4n+6} + \frac{s^{2n+12}}{2n+12} \right] \end{aligned}$$

$$\begin{aligned} NH(W_n : S, t) &= 2n \left[ \frac{1}{4n+6} + \frac{1}{2n+12} \right] \\ &= \frac{3n(n+3)}{(n+6)(2n+3)}. \end{aligned}$$

$$(6) \quad NI(W_n : S, t) = S_s J D_s D_t [NM(W_n : s, t)]_{s=t=1}$$

$$D_s D_t [NM(W_n)] = 3n^2(n+6)s^{n+6}t^{3n} + n(n+6)^2s^{n+6}t^{n+6}$$

$$J(D_s D_t) = 3n^2(n+6)s^{4n+6} + n(n+6)^2s^{2n+12}$$

$$\begin{aligned} S_x J(D_s D_t) &= \int_0^s 3n^2(n+6)x^{4n+6} + n(n+6)^2x^{2n+11} dx \\ &= 3n^2(n+6) \frac{s^{4n+6}}{4n+6} + n(n+6)^2 \frac{s^{2n+12}}{2n+12} \end{aligned}$$

$$\begin{aligned} NI(W_n : S, t) &= 3n^2(n+6) \frac{1}{4n+6} + n(n+6)^2 \frac{1}{2n+12} \\ &= n(n+6) \left[ \frac{3n}{4n+6} + \frac{n+6}{2(n+6)} \right] \\ &= \frac{n(n+6)(5n+3)}{2(2n+3)}. \end{aligned}$$

□

**2.2. Helm graph.** The helm graph is the graph obtained from a wheel graph by adjoining a pendant edge at each node of the cycle.

Let  $H_n$  be the helm graph having number of edges  $3n$  and number of vertices  $2n+1$ . The neighborhood degree sequence of edge partition of  $H_n$  is given below

$$(2.2) \quad E_{(9+n)(4)} = n, \quad E_{(9+n)(9+n)} = n, \quad E_{(9+n)(9+n)} = n$$

**Theorem 2.3.** *NM-polynomial of helm graph is given by,*

$$NM(H_n : s, t) = ns^{n+9} [t^{4n} + t^4 + t^{n+9}].$$

*Proof.* By using equation (2.2)

$$\begin{aligned} NM(H_n : s, t) &= m_{(n+9)4n} s^{n+9} t^{4n} + m_{(n+9)4} s^{n+9} t^4 + m_{(n+9)(n+9)} s^{n+9} t^{n+9} \\ &= ns^{n+9} t^{4n} + ns^{n+9} t^4 + ns^{n+9} t^{n+9} \\ &= ns^{n+9} [t^{4n} + t^4 + t^{n+9}]. \end{aligned}$$

□

**Theorem 2.4.** *Let  $H_n$  be the helm graph, then*

$$(1) \text{ Neighborhood First Zagreb index, } M'_1(H_n : s, t) = 8n(n+5).$$

$$(2) \text{ Neighborhood Second Zagreb index, } M_2^*(H_n : s, t) = n(5n^2 + 58n + 117).$$

$$(3) \text{ Neighborhood Modified Zagreb index, } {}^{nm}M_2(H_n : s, t) = \frac{n^2 + 14n + 9}{4(n+9)^2}.$$

$$(4) \text{ Neighborhood Symmetric division deg index, } NSDD(H_n : s, t) = \frac{n^3 + 23n^2 + 155n + 133}{4(n+9)}.$$

$$(5) \text{ Neighborhood Harmonic Index, } NH(H_n : s, t) = \frac{n(17n^2 + 226n + 513)}{(5n+9)(n+13)(n+9)}.$$

$$(6) \text{ Neighborhood Inverse Index } NI(H_n : s, t) = \frac{13n^3 + 335n^2 + 2151n + 1701}{2(5n+9)(n+13)}.$$

*Proof.* Using equation (2.2) and Table 1

$$(1) D_s [NM(H_n)] = s \frac{\partial}{\partial s} [NM(H_n)] = sn(n+9)s^{n+8} [t^{4n} + t^4 + t^{n+9}].$$

$$= n(n+9)s^{n+9} [t^{4n} + t^4 + t^{n+9}]$$

$$D_t [NM(H_n)] = t \frac{\partial}{\partial t} [NM(H_n)] = tns^{n+9} [4nt^{4n-1} + 4t^3 + (n+9)t^{n+8}]$$

$$= ns^{n+9} [4nt^{4n} + 4t^4 + (n+9)t^{n+9}]$$

$$M_1'(H_n) = (D_s + D_t) [NM(H_n; s, t)]_{s=t=1}$$

$$= 8n(n+5).$$

$$(2) D_s D_t [NM(H_n)] = s \frac{\partial}{\partial s} [D_t] = sn(n+9)s^{n+8} [4nt^{4n} + 4t^4 + (n+9)t^{n+9}]$$

$$= n(n+9)s^{n+9} [4nt^{4n} + 4t^4 + (n+9)t^{n+9}]$$

$$M_2^*(H_n) = (D_s D_t) [NM(H_n; s, t)]_{s=t=1} = n(n+9)[4n+4+(n+9)]$$

$$= 5n^3 + 58n^2 + 117.$$

$$(3) S_s S_t = \int_0^s \frac{S_t(x, t)}{x} dx$$

$$= n \frac{s^{n+9}}{n+9} \left[ \frac{t^{4n}}{4n} + \frac{t^4}{4} + \frac{t^{n+9}}{n+9} \right]$$

$${}^m M_2(H_n : s, t) = (S_s S_t) [NM(H_n; s, t)]_{s=t=1}$$

$$= \frac{n}{n+9} \left[ \frac{1}{4n} + \frac{1}{4} + \frac{1}{n+9} \right]$$

$$= \frac{n^2 + 14n + 9}{4(n+9)^2}.$$

$$\begin{aligned}
 (4) \quad S_s D_t &= \int_0^s \frac{D_t(x, t)}{x} dx \\
 &= \int_0^s n x^{n+8} [4nt^{4n} + 4t^4 + (n+9)t^{n+9}] dx \\
 &= n \frac{s^{n+9}}{n+9} [4nt^{4n} + 4t^4 + (n+9)t^{n+9}]
 \end{aligned}$$

$$\begin{aligned}
 D_s S_t &= s \frac{\partial}{\partial s} (S_t) \\
 &= s \frac{\partial}{\partial s} \left[ n s^{n+9} \left( \frac{t^{4n}}{4n} + \frac{t^4}{4} + \frac{t^{n+9}}{n+9} \right) \right] \\
 &= n(n+9) s^{n+9} \left[ \frac{t^{4n}}{4n} + \frac{t^4}{4} + \frac{t^{n+9}}{n+9} \right]
 \end{aligned}$$

$$\begin{aligned}
 NSDD(H_n : s, t) &= [D_s S_t + S_s D_t] [NM(H_n; s, t)]_{s=t=1} \\
 &= n(n+9) \left[ \frac{1}{4n} + \frac{1}{4} + \frac{1}{n+9} \right] + \frac{n}{n+9} [4n + 4n + n + 9] \\
 NSDD(H_n) &= \frac{n^3 + 43n^2 + 187n + 81}{4(n+9)}.
 \end{aligned}$$

$$\begin{aligned}
 (5) \quad J[NM(H_n; s, t)] &= NM(H_n; s, s) \\
 &= n s^{n+9} [s^{4n} + s^4 + s^{n+9}] \\
 &= n [s^{5n+9} + s^{n+13} + s^{2n+18}]
 \end{aligned}$$

$$\begin{aligned}
 S_s J[NM(H_n)] &= \int_0^s \frac{J[NM(H_n)]_x}{x} dx \\
 &= n \int_0^s (x^{5n+8} + x^{n+12} + x^{2n+17}) dx \\
 &= n \left[ \frac{s^{5n+9}}{5n+9} + \frac{s^{n+13}}{n+13} + \frac{s^{2n+18}}{2n+18} \right]
 \end{aligned}$$

$$\begin{aligned}
 NH(H_n : S, t) &= 2 S_s J[NM(H_n : s)]_{s=1} \\
 &= 2n \left[ \frac{1}{5n+9} + \frac{1}{n+13} + \frac{1}{2n+18} \right]
 \end{aligned}$$

$$NH(H_n : S, t) = \frac{n(17n^2 + 226n + 513)}{(5n+9)(n+13)(n+9)}.$$

$$(6) \quad NI(H_n : S, t) = S_s J D_s D_t \left[ NM(H_n : s, t) \right]_{s=t=1}$$

$$D_s D_t [NM(H_n)] = n(n+9)s^{n+9} \left[ 4nt^{4n} + 4t^4 + (n+9)t^{n+9} \right]$$

$$J(D_s D_t) = (n+9)s^{n+9} \left[ 4ns^{4n} + 4s^4 + (n+9)s^{n+9} \right]$$

$$= (n+9) \left[ 4ns^{5n+9} + 4s^{n+13} + (n+9)s^{2n+18} \right]$$

$$S_s J(D_s D_t) = \int_0^s (n+9)s^{n+9} \left[ 4ns^{4n} + 4s^4 + (n+9)s^{n+9} \right] dx$$

$$= (n+9) \left[ 4n \frac{s^{5n+9}}{5n+9} + 4 \frac{s^{n+13}}{n+13} + (n+9) \frac{s^{2n+18}}{2n+18} \right]$$

$$NI(H_n : S, t) = (n+9) \left[ \frac{4n}{5n+9} + \frac{4}{n+13} + \frac{(n+9)}{2n+18} \right]$$

$$NI(H_n : s, t) = \frac{13n^3 + 335n^2 + 2151n + 1701}{2(5n+9)(n+13)}.$$

□

**2.3. Gear graph.** A Gear graph is obtained by adding a vertex to each outer edge of Wheel graph.

Let  $G_n$  be the gear graph with  $3n$  number of edges and  $2n+1$  number of vertices and its edge partition of neighbourhood degree sequence of  $G_n$  is

$$(2.3) \quad E_{(n+4)(3n)} = n, \quad E_{(n+4)(6)} = 2n$$

**Theorem 2.5.** *NM-polynomial of gear graph is given by,*

$$NM(G_n : s, t) = ns^{n+4} \left[ t^{3n} + 2t^6 \right].$$

*Proof.* By using the equation (2.3)

$$\begin{aligned} NM(G_n : s, t) &= m_{(n+4)(3n)} s^{n+4} t^{3n} + m_{(n+4)(6)} s^{n+4} t^6 \\ &= n s^{n+4} t^{3n} + 2n s^{n+4} t^6 \\ &= ns^{n+4} \left[ t^{3n} + 2t^6 \right]. \end{aligned}$$

□

**Theorem 2.6.** *Let  $G_n$  be the gear graph, then*

$$(1) \quad \text{Neighborhood First Zagreb index, } M'_1(G_n : s, t) = 6n(n+4).$$

$$(2) \quad \text{Neighborhood Second Zagreb index, } M_2^*(G_n : s, t) = 3n(n+4)^2.$$

$$(3) \quad \text{Neighborhood Modified Second Zagreb index, } {}^{nm}M_2(G_n : s, t) = \frac{n+1}{n+4}.$$

$$(4) \text{ Neighborhood Symmetric division deg index, } NSDD(G_n : s, t) = \frac{n^2 + 14n + 4}{3}.$$

$$(5) \text{ Neighborhood Harmonic Index, } NH(G_n : s, t) = \frac{9n(n+2)}{2(n+1)(n+10)}.$$

$$(6) \text{ Neighborhood Inverse Index, } NI(G_n : s, t) = \frac{3n(n+4)(n^2 + 26n + 16)}{4(n+1)(n+10)}.$$

*Proof.* Using equation (2.3) and Table 1

$$\begin{aligned} (1) \quad D_s [NM(G_n)] &= s \frac{\partial}{\partial s} [NM(G_n)] = sn(n+4)s^{n+3} [t^{3n} + 2t^6] \\ &= n(n+4)s^{n+4} [t^{3n} + 2t^6] \end{aligned}$$

$$\begin{aligned} D_t [NM(G_n)] &= t \frac{\partial}{\partial t} [NM(G_n)] = t n s^{n+4} (3nt^{3n-1} + 12t^5) \\ &= n s^{n+4} (3nt^{3n} + 12t^6) \end{aligned}$$

$$\begin{aligned} M'_1(G_n) &= (D_s + D_t) [NM(G_n; s, t)]_{s=t=1} \\ &= 6n(n+4). \end{aligned}$$

$$\begin{aligned} (2) \quad D_s D_t [NM(G_n)] &= s \frac{\partial}{\partial s} (D_t) = s n (3nt^{3n} + 12t^6)(n+4)s^{n+3} \\ &= n(n+4)s^{n+4} (3nt^{3n} + 12t^6) \end{aligned}$$

$$\begin{aligned} M_2^*(G_n) &= (D_s D_t) [NM(G_n; s, t)]_{s=t=1} = n(n+4)(3n+12) \\ &= 3n(n+4)^2. \end{aligned}$$

$$\begin{aligned} (3) \quad S_s S_t &= \int_0^s \frac{S_t(x, t)}{x} dx \\ &= \int_0^s nx^{n+3} \left[ \frac{t^{3n}}{3n} + \frac{t^6}{3} \right] dx \\ &= n \frac{s^{n+4}}{n+4} \left[ \frac{t^{3n}}{3n} + \frac{t^6}{3} \right] \end{aligned}$$

$$\begin{aligned} {}^{nm}M_2(G_n : s, t) &= (S_s S_t) [NM(H_n; s, t)]_{s=t=1} \\ &= \frac{n}{n+4} \left[ \frac{1}{3n} + \frac{1}{3} \right] \\ &= \frac{n+1}{n+4}. \end{aligned}$$

$$\begin{aligned} (4) \quad S_s D_t &= \int_0^s \frac{D_t(x, t)}{x} dx \\ &= \int_0^s nx^{n+3} [3nt^{3n} + 12t^6] dx \end{aligned}$$

$$= 3n \frac{s^{n+4}}{n+4} [nt^{3n} + 4t^6]$$

$$D_s S_t = s \frac{\partial}{\partial s} (S_t)$$

$$= s \frac{\partial}{\partial s} \left[ ns^{n+4} \left( \frac{t^{3n}}{3n} + \frac{t^6}{3} \right) \right]$$

$$= n(n+4)s^{n+4} \left[ \frac{t^{3n}}{3n} + \frac{t^6}{3} \right]$$

$$\left[ D_s S_t + S_s D_t \right]_{s=t=1} = n(n+4) \left[ \frac{1}{3n} + \frac{1}{3} + \frac{3}{n+4} \right]$$

$$NSDD(G_n : s, t) = \left[ D_s S_t + S_s D_t \right] \left[ NM(G_n; s, t) \right]_{s=t=1}$$

$$NSDD(G_n) = \frac{1}{3}(n^2 + 14n + 4).$$

$$(5) \quad J[NM(G_n; s, t)] = NM(G_n; s, s)$$

$$= ns^{n+4} [s^{3n} + 2s^6]$$

$$= n [s^{4n+4} + 2s^{n+10}]$$

$$S_s J[NM(G_n)] = \int_0^s \frac{J[NM(G_n)]_x}{x} dx$$

$$= n \int_0^s (x^{4n+3} + 2x^{n+9}) dx$$

$$= n \left[ \frac{s^{4n+4}}{4n+4} + \frac{2s^{n+10}}{n+10} \right]$$

$$NH(G_n : S, t) = 2S_s J[NM(G_n : s)]_{s=1}$$

$$= 2n \left[ \frac{1}{4n+4} + \frac{2}{n+10} \right]$$

$$NH(G_n : S, t) = \frac{9n(n+2)}{2(n+1)(n+10)}.$$

$$(6) \quad D_s D_t [NM(G_n)] = n(n+4)s^{n+4} [3nt^{3n} + 12t^6]$$

$$J(D_s D_t) = n(n+4)s^{n+4} [3ns^{3n} + 12s^6]$$

$$= 3n^2(n+4)s^{4n+4} + 12n(n+4)s^{n+10}$$

$$S_s J(D_s D_t) = \int_0^s \left[ 3n^2(n+4)x^{4n+3} + 12n(n+4)x^{n+9} \right] dx$$

$$= 3n^2(n+4) \frac{s^{4n+4}}{4n+4} + 12n(n+4) \frac{s^{n+10}}{n+10}$$

$$NI(G_n : S, t) = \frac{3n^2(n+4)}{4n+4} + \frac{12n(n+4)}{n+10}$$

$$NI(G_n : s, t) = \frac{3n(n+4)(n^2 + 26n + 16)}{4(n+1)(n+10)}.$$

□

### 3. COMPARISON

In all three graphs shown in Figures 1-3,  $NSDD$  shows the most substantial and consistent growth, maintaining a steady linear increase without slowing down, but  $M_1'$ ,  $M_2$ , and  $M_2^*$  grow exponentially, with  $M_2^*$  showing the most rapid growth.  $NH$  grows rapidly at first but begins to plateau, suggesting that its growth has an upper limit or reaches a steady state.  $NI$  grows very slowly in comparison, with a much smaller increase over the same range of  $n$ .  $nmM_2$  is nearly flat, indicating a stable, unchanging value throughout the range, suggesting that it is unaffected by the increase in  $n$ .




FIGURE 1. Wheel graph




FIGURE 2. Helm graph




FIGURE 3. Gear graph

#### 4. CONCLUSION

The NM-polynomial and related topological indices derived for cycle-based graphs such as wheel, helm, and gear graphs have broad applications in theoretical chemistry and material science. These descriptors are instrumental in QSAR/QSPR modeling, molecular database indexing, and predicting physicochemical properties of ring-structured compounds. Additionally, they offer valuable insights into structural complexity and connectivity in network science and nanostructure design, demonstrating the wide-reaching utility of the methods developed in this study.

A promising future direction is to extend these methods to dynamic and weighted networks, such as biological interaction networks or evolving chemical reaction networks. Incorporating temporal or weighted connectivity into NM-polynomials could provide deeper insights into time-dependent properties, enabling better modeling of dynamic molecular systems, smart materials, or responsive nanostructures. This would broaden the applicability from static structures to real-world, evolving systems in both chemistry and materials science.

#### REFERENCES

- [1] K.C. Das, K. Xu and J. Nam, *On Zagreb indices of graphs*, Front. Math. China, 10(3), (2015), 567-582.
- [2] E. Deutsch and S. Klavzar, *M-polynomial and degree-based topological indices*, Iran. J. Math. Chem., 6, (2015), 93-102.
- [3] J. Devillers and A. Balaban, **Topological indices and related descriptors in QSAR and QSPR**, Taylor and Francis, CRC Press, 1st edition, (1999).

- [4] Deepasree S. K, Ranjini P.S, V. Lokesha, Phaniraju, *Topological indices of some operations in cycle graphs related to EV and VE indices*, Montes Taurus J. Pure Appl. Math6(1)1–11, 2024.
- [5] Deepasree S. Kumar, P. S. Ranjini, V. Lokesha, Dafik, I. H. Agustin; NV L index and bounds for the operations of graphs. AIP Conf. Proc. 30 July 2024; 3176 (1): 020012.
- [6] V. R. Kulli, *Neighborhood Indices of Nanostructures*, International Journal of Current Research in Science and Technology, 5(3), (2019), 1-14.
- [7] V. Lokesha and S. Jain, *New results on the F-index of graphs based on a corona type products of graphs*, Proc. Jangjeon Math. Soc., 23(2), (2020), 141-148.
- [8] V. Lokesha, Suvarna, A. S. Cevik and I. N. Cangul, *VL Reciprocal Status Index and Co-Index of Graphs*, Journal of Mathematics, Article ID 5529080, 10 pages, 2021.
- [9] V. Lokesha and Suvarna *Bounds for VL Status Index and Coindex of Graphs and Validate to Few Specific Chapter 20*, Applied Nonlinear Analysis and Soft Computing, Advances in Intelligent Systems and Computing 1437, Springer Nature Singapore Pte Ltd. H. Dutta et al. (eds.), (2023).
- [10] V. Lokesha, Suvarana, A. Sinan Cevik, *V L status index and co-index of connected graphs*, Proc of the Jangjeon Mathematical society, 24(3) (2021), 285-295.
- [11] Lokesha, V, Maragadam A.S, Suvarna Cangul, I.N, *Topological coindices of phytochemicals examined for COVID-19 therapy*, Proc. of the Jangjeon Mathematical society, 26(4), 2023 381-405.
- [12] V. Lokesha, Suvarna, M. Manjunath and K. Zeba Yasmeen, *VL Temperture index of certain archimedean lattice*, South East Asian J. of Mathematics and Mathematical Sciences, 17(1), (2021), 213-222.
- [13] S. Mondal, N. De and A. Pal, *On Neighbourhood Zagreb index of product graphs*, J. Mol. Struct., (2021).
- [14] S. Mondal, N. De and A. Pal, *On some new neighbourhood degree based indices*, Acta Chemica Iasi, 27, (2019), 31-46.
- [15] S. Mondal, N. De and A. Pal, *Topological Properties of Networks Using M-Polynomial Approach*. Konuralp Journal of Mathematics, 8, (2020), 97-105.
- [16] S. Mondal, M. K. Siddiqui, N. De and A. Pal, *Neighbourhood M-polynomial of crystallographic structures*, Biointerface Research in Applied Chemistry, 11(2), (2021), 9372-9381.
- [17] S. Mondal, M. Imran, N. De and A. Pal, *Neighbourhood M-polynomial of titanium compounds*. Arabian Journal of Chemistry, (2021).
- [18] Z. Shao; P. Wu, Y. Gao, I. Gutman and X. Zhang, *On the maximum ABC index of graphs without pendent vertices*, Appl. Math. Comput., 315, (2017), 298-312.
- [19] Z. Shao, P. Wu, X. Zhang, D. Dimitrov and J. Liu, *On the maximum ABC index of graphs with prescribed size and without pendent vertices*, IEEE Access 6, (2018), 27604-27616.
- [20] A. M. Soner, N. D. Naji, and I. Gutman, *On leap Zagreb indices of graphs*, Commun. Comb. Optim., 2, (2017), 99-117.
- [21] Suvarna, V. Lokesha, Shantha Kumari *On VL Recciproal Status Index And Co-Index Of Connected Graphs-II*, Proc of the Jangjeon Mathematical society, 25(3) (2022), 319-329.
- [22] D. Vukicevic and M. Gasperov, *Bond additive modeling 1. Adriatic indices*, Croatica chemica acta, 83(3), (2010), 243-260.
- [23] Wiener H, *Structural determination of paraffin boiling points*, J.Am.Chem.Soc., 69, (1947), 17-20.