

ON THE GENERALIZATION OF THE TORSION FUNCTOR AND P -SEMIPRIME MODULES OVER NONCOMMUTATIVE RINGS

Teklemichael Worku Bihonegn , Tilahun Abebaw and Nega Arega

ABSTRACT. Let R be an associative Noetherian unital noncommutative ring R . We introduce the functor $P\Gamma_P$ over the category of R -modules and use it to characterize P -semiprime. P -semisecond R -modules also characterized by the functor $P\Lambda_P$. We also show that the Greenless-May type Duality (GM) and Matlis Greenless-May Equality (MGM) hold over the full subcategory of R -Mod consisting of P -semiprime and P -semisecond modules. Finally, we generate a one-sided right ideal $P\Gamma_P(R)$, which gives an equivalent formulation to solve Köthe conjecture positively or negatively.

Key Words: P -semiprime, P -semisecond, torsion functor, adic completion and Köthe conjecture.

2010 Mathematics Subject Classification: Primary: 16D90; Secondary: 16N60.

1. INTRODUCTION

Let R be a unital Noetherian ring which is not necessarily commutative. R is semiprime if for all ideals P of R , $P^2 = 0$ implies that $P = 0$. R is reduced if for all $a \in R$, $a^2 = 0$ implies that $a = 0$. If R is commutative, the two notions coincides. An ideal P of a ring R is semiprime (resp. completely semiprime) if the quotient ring R/I is a semiprime

Received: 04 January 2024, Accepted: 15 May 2024. Communicated by Alireza Naghipour;
*Address correspondence to T. W. Bihonegn; E-mail: teklemichael.worku@aaau.edu.et.

This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/).

Copyright © 2024 The Author(s). Published by University of Mohaghegh Ardabili.

(resp. reduced) ring. Any reduced ring is semiprime. However, the ring $M_2(\mathbb{Z})$ is a semiprime ring which is not reduced. The notion of reduced, semiprime, semisecund and coreduced modules has been widely studied see for instance [11, 15, 16, 18, 21]. When R is commutative, the locally nilradical functor $a\Gamma_a(-)$ over a category of R -modules, where a is the ring element has been studied as a measure of how far a module from being reduced in [9]. Moreover, in [19], it has been studied about P -reduced and P -coreduced modules in relation with Matlis-Greenlees-May Equivalence and Greenlees-May duality. The Köthe conjecture states that if a ring R has no nonzero nil ideals then R has no nonzero nil one-sided ideals and it has existed since 1930. Even if the problem is still open lots of equivalent formulations has been made. The sum of two right nil ideals in any ring R is nil is also an equivalent formulation for the conjecture, [1, 8, 14]. This paper is organized as follows: In Section 2 we introduce the functor $P\Gamma_P$ and show that it is a 1) radical over a category of R -modules; 2) left exact over an abelian full subcategory of $R\text{-Mod}$ consisting of flat modules; 3) we also use it to characterize the P -semiprime modules; 4) we characterize P -semisecund modules using the functor $P\Lambda_P$. In Section 3 we study applications of P -semiprime and P -semisecund modules and show that the Greenlees-May duality and Matlis-Greenlees-May Equality holds. In Section 4, we generate a right nil ideal by considering $P\Gamma_P$ over rings. We also provide a gadget that produces one sided nil ideals for any noncommutative ring for a given right ideal (Proposition 4.2). This is useful in the possible counter examples to answer the Köthe conjecture in the negative. Finally, we pose questions which are equivalent to solve the Köthe conjecture in the negative (Question 4.8) and positive (Question 4.9) using the ideal $P\Gamma_P$.

2. P -SEMPIRIME AND P -SEMISECOND MODULES

In this section unless otherwise mentioned $R\text{-Mod}$ represents the category of left R -modules. Let P be an ideal of a ring R . A submodule N of an R -module M is P -semiprime if for all $m \in M$, $P^2m \subseteq N$ implies that $Pm \subseteq N$. A submodule N of an R -module M is semiprime if it is P -semiprime for all ideals P of R . An R -module M/N is semiprime (resp. P -semiprime) if N is a semiprime (resp. P -semiprime) submodule of M .

Remark 2.1. A ring R is semiprime if and only if the R -module R is semiprime.

Let $R\text{-Mod}$ be a category of left R -modules. A functor $\gamma(-) : R\text{-Mod} \rightarrow R\text{-Mod}$ is a *preradical* if $\gamma(M)$ is a submodule of M and for every R -homomorphism $f : M \rightarrow N$, $f(\gamma(M)) \subseteq \gamma(N)$. γ is a *radical* if it is a preradical and for all $M \in R\text{-Mod}$, $\gamma(M/\gamma(M)) = 0$. A radical γ is left exact if for every submodule N of a module $M \in R\text{-Mod}$, $\gamma(N) = N \cap \gamma(M)$. Equivalently, if for any exact sequence $0 \rightarrow N \rightarrow M \rightarrow K$ of R -modules, the sequence $0 \rightarrow \gamma(N) \rightarrow \gamma(M) \rightarrow \gamma(K)$ is also exact.

Definition 2.2. Let R be a Noetherian ring and P a right ideal of R . $\Gamma_P(-) : R\text{-Mod} \rightarrow R\text{-Mod}$ $M \mapsto \Gamma_P(M) := \{m \in M : P^k m = 0, \text{ for some } k \in \mathbb{Z}^+\}$.

Proposition 2.3. *The functor $\Gamma_P(-) : R\text{-Mod} \rightarrow R\text{-Mod}$ is a left exact radical.*

Proof. 1. Consider the R -module homomorphism $f : M \rightarrow N$. Let $y \in f(\Gamma_P(M))$, $y = f(m) \in N$ for some $m \in \Gamma_P(M)$, i.e., $P^k m = 0$ for some positive integer k . Now, $P^k y = P^k f(m) = f(P^k m) = f(0) = 0$ which implies $y \in \Gamma_P(N)$.

2. To show it is radical, it is enough to show that $\Gamma_P(M/\Gamma_P(M)) = 0$. Let $y \in \Gamma_P(M/\Gamma_P(M))$ such that $P^k m \in \Gamma_P(M)$, where $y = m + \Gamma_P(M)$ for some $m \in M$. Then there exists a positive integer k_1 such that $P^{k_1}(P^k m) = 0$ which implies $P^{k_1+k} m = 0$. It follows that $m \in \Gamma_P(M)$ and thus $y = 0$.

3. It is similar to the proof of [3, Lemma 1.16] □

By multiplying the torsion functor Γ_P by P from the left we define $P\Gamma_P$ as follows:

Definition 2.4. Let P be an ideal of a ring R . A functor

$P\Gamma_P(-) : R\text{-Mod} \rightarrow R\text{-Mod}$ is defined by

$$M \mapsto P\Gamma_P(M) := \left\{ \sum_{i=1}^n r_i m_i : r_i \in P \text{ and } m_i \in \Gamma_P(M) \right\}.$$

Proposition 2.5. *Let $M \in R\text{-Mod}$ and P an ideal of R . The following are equivalent:*

1. M is P -semiprime.
2. $(0 : m)$ is an P -semiprime left ideal of R for all $0 \neq m \in M$.
3. $(0 :_M P) = (0 :_M P^2)$.
4. $\text{Hom}_R(R/P, M) = \text{Hom}_R(R/P^2, M)$.

5. $\Gamma_P(M) \cong \text{Hom}(R/P, M)$.
6. $P\Gamma_P(M) = 0$.

Proof. $1 \Rightarrow 2$ For any left ideal P of R , let $P^2 \subseteq (0 : m)$. Then this implies that $P^2m = 0$ for all nonzero $m \in M$, since M is P -semiprime R -module it follows that $Pm = 0$ and hence $P \subseteq (0 : m)$.

$2 \Rightarrow 1$ For any ideal P of R and $0 \neq m \in M$, let $P^2m = 0$ implies $P^2 \in (0 : m)$ then by hypothesis $P \in (0 : m)$ which implies $Pm = 0$, thus P -semiprime.

$2 \Rightarrow (3)$ Since $(0 : m)$ is P -semiprime ideal of R , it follows that $(0 :_M P^2) \subseteq (0 :_M P)$, the other inclusion is obvious.

$3 \Rightarrow 4$ Since $(0 :_M P)$ is a left R -module, then it coincides with $\text{Hom}_R(R/P, M)$ then the result follows.

$(4) \Rightarrow (5)$ since $\Gamma_P(M) \cong \varinjlim_k \text{Hom}_R(R/P^k, M)$. then by (4) we

have

$\text{Hom}_R(R/P, M) = \text{Hom}_R(R/P^k, M)$ for all $k \in \mathbb{Z}^+$. So, $\Gamma_P(M) \cong \text{Hom}_R(R/P, M)$.

$(5) \Rightarrow (6)$ $P\Gamma_P(M) \cong P(\text{Hom}(R/P, M)) = 0$.

$1 \Rightarrow 6$ Let M be P -semiprime module, $m \in P\Gamma_P(M)$ and $m = \sum_{i=1}^n a_i m_i$, $a_i \in P$ and $m_i \in \Gamma_P(M)$, i.e., $P^{k_i} m_i = 0$ for some positive integers k_i . By hypothesis $Pm_i = 0$ then for each $a_i \in P$ we have $a_i m_i = 0$ then $m = 0$. So, $P\Gamma_P(M) = 0$.

$6 \Rightarrow 1$ Suppose $P\Gamma_P(M) = 0$. Let $m \in M$ and $P^2m = 0$ which implies $m \in \Gamma_P(M)$ then $Pm \subseteq P\Gamma_P(M) = 0$, so M is P -semiprime R -module. □

So, the functor $P\Gamma_P(-)$ on $R\text{-Mod}$ is a measure of how far a module is from being P -semiprime.

A left R -module F is *flat* if the functor $- \otimes F$ is exact. A short exact sequence $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ of left R -modules is called *pure* if $K \otimes L \rightarrow K \otimes M$ is a monomorphism for every right R -module K .

In general, $P\Gamma_P$ is not left exact. Consider the \mathbb{Z} -module $M = \mathbb{Z}_{12}$ and the submodule $N = \{0, 6\}$ and take $I = \langle 2 \rangle$. Now, $\langle 2 \rangle \Gamma_{\langle 2 \rangle}(M) = \{0, 3, 6, 9\}$, $\langle 2 \rangle \Gamma_{\langle 2 \rangle}(N) = \{0\}$ and $\langle 2 \rangle \Gamma_{\langle 2 \rangle}(M) \cap N = N$, but $\langle 2 \rangle \Gamma_{\langle 2 \rangle}(N) = \{0\}$. Let $\text{Fl}(R)$ denote an abelian full subcategory of $R\text{-Mod}$ consisting of all flat R -modules. This abelian category was studied in [6, 7, 20]. Note that since any free R -module is flat, $R \in \text{Fl}(R)$.

Proposition 2.6. Consider the functor

$$P\Gamma_P(-) : R\text{-Mod} \rightarrow R\text{-Mod}, M \mapsto P\Gamma_P(M).$$

Then,

1. $P\Gamma_P(-)$ is a radical and it is left exact over $\text{Fl}(R)$.
2. $P\Gamma_P(M) \cong P \otimes_R \Gamma_P(M)$, i.e., a composition of $\Gamma_P(-)$ and $P \otimes -$.
3. If in addition $\text{Fl}(R)$ has enough injectives, then $P\Gamma_P$ is exact.

Proof. 1. To show $P\Gamma_P$ is a preradical, consider the module homomorphism $f : M \rightarrow N$ for any left R -modules M and N . Let $y \in f(P\Gamma_P(M))$, $y = f(m) \in N$, $m = \sum_{i=1}^l r_i m_i$ and $r_i \in P, m_i \in \Gamma_P(M)$, i.e., $P^k m_i = 0$ for some positive integer k . Now, $P^k(y) = P^k(f(m)) = f(P^k \sum_{i=1}^l r_i m_i) = f(0) = 0$ which implies $y \in \Gamma_P(N)$. To show it is radical, it is enough to show that $M/P\Gamma_P(M)$ is P -semiprime. Let $\bar{m} \in M/P\Gamma_P(M)$ and $P^2 \bar{m} = \bar{0}$ which implies $P^2(m + P\Gamma_P(M)) = P\Gamma_P(M)$, then $P^2 m \in P\Gamma_P(M)$ and hence $Pm \in \Gamma_P(M)$ such that $P^k m = 0$ for some positive integer k which implies $m \in \Gamma_P(M)$ and so, $Pm \subseteq P\Gamma_P(M)$ and thus, $P\bar{m} = \bar{0}$. Then by Proposition 2.5 the functor is radical.

$P\Gamma_P$ is left exact if and only if for all submodules N of M $P\Gamma_P(N) = P\Gamma_P(M) \cap N$, [12]. Since, $P\Gamma_P(N)$ is submodule of both $P\Gamma_P(M)$ and N it is easy to show that $P\Gamma_P(N) \subseteq P\Gamma_P(M) \cap N$. However, by hypothesis every submodule is pure, so $P\Gamma_P(N) = PM \cap \Gamma_P(N)$, [5, Proposition 8.1]. Then it follows that $PM \cap \Gamma_P(N) \subseteq P\Gamma_P(M) \cap N$. To show the reverse inclusion let $y \in P\Gamma_P(M) \cap N$. Then, $y = \sum_{i=1}^n r_i m_i$ such that $P^{k_i} r_i = 0$ for some positive integers k_i , $1 \leq i \leq n$. Now, $P^k y = \sum_{i=1}^n P^k r_i m_i = 0$, where $k = k_1 + \dots + k_n$, which shows that $y \in PM \cap \Gamma_P(N)$. Hence, $P\Gamma_P(N) = P\Gamma_P(M) \cap N$.

2. Since $M \in \text{Fl}(R)$ and $\text{Fl}(R)$ is an abelian subcategory, $\Gamma_P(M) \in \text{Fl}(R)$. It follows that $P\Gamma_P(M) \cong P \otimes_R \Gamma_P(M)$.
3. Since $P\Gamma_P$ is left exact by part 1 and $\text{Fl}(R)$ has enough injectives, we can compute the right derived functor of $P\Gamma_P$. By [17, Theorem 10.47] $\mathbf{R}^i(P\Gamma_P(M)) \cong \mathbf{R}^i(P \otimes_R \Gamma_P(M)) \cong \mathbf{R}^i(P \otimes_R \mathbf{R}^i(\Gamma_P(M))) \cong 0$. It is zero because $P \otimes_R -$ is exact. \square

In [4] examples for which $\text{Fl}(R)$ has enough injectives were given. This happens when R_P is quasi-Frobenius for all $P \in \text{ASS}(R)$, the

assassinator of R . However, the only two examples of rings which were given namely; commutative Noetherian domain and $R = k[x, y]/\langle xy \rangle$ are both reduced and flat modules over reduced rings in which case, our functor $P\Gamma_P(-) : \text{Fl}(R) \rightarrow \text{Fl}(R)$ will be trivial. see [4, Theorem 3].

Example 2.7. If $R = k[t]/\langle t^2 \rangle$. Then R is a flat R -Mod which is not semiprime and also $\langle t \rangle \Gamma_{\langle t \rangle}(R) \neq 0$.

Definition 2.8. Let P be an ideal of R . Define the P -adic completion functor $\Lambda_P(-) : R\text{-Mod} \rightarrow R\text{-Mod}$ by $M \mapsto \Lambda_P(M) := \varprojlim_k M/P^k M$.

Definition 2.9. Let P be an ideal of R . A left R -module M is said to be P -semisecund if $P^2 M = PM$.

A left R -module M is said to be semisecund if M is P -semisecund module for every ideal P of R , [2].

Proposition 2.10. For any ideal P of R and R -module M the following are equivalent:

1. M is P -semisecund.
2. $R/P \otimes_R M \cong R/P^2 \otimes_R M$,
3. $R/P \otimes M \cong \Lambda_P(M)$.
4. $P\Lambda_P(M) = 0$.

Proof. (1) \Rightarrow (2) $R/P \otimes_R M \cong M/PM$ since M is semisecund $R/P \otimes_R M \cong M/P^2 M \cong R/P^2 \otimes_R M$.

$$(2) \Rightarrow (3) \Lambda_P(M) = \varprojlim_k (M/P^k M) \cong \varprojlim_k (M/P^k \otimes_R M) \cong$$

$$\varprojlim_k (M/P \otimes_R M) = R/P \otimes_R M.$$

$$(3) \Rightarrow (4) P\Lambda_P(M) \cong P(R/P \otimes_R M) = P(M/PM) = 0$$

(1) \Rightarrow (4) Since $P^2 M = PM$, it follows that $P^k M = PM$ for each positive integer k , then $P\Lambda_P(M) = 0$

(4) \Rightarrow (1) Since $P\Lambda_P(M) = 0$ which implies $P(\varprojlim_k M/P^k M) = 0$,

then $PM = P^k M$ for all $k \in \mathbb{Z}^+$ which implies $PM = P^2 M$. \square

3. APPLICATIONS OF P -SEMIPRIME AND P -SEMISECOND MODULES

If R is commutative ring and P is an ideal of R , then P -reduced and P -semiprime coincide and similarly, P -coreduced also coincide with P -semisecond modules. So, Matlis-Greenlees-May Equivalence and Greenlees-May type duality holds, see [19]. However, when R is noncommutative the above notions are different. Hence in this section we prove that Matlis-Greenlees-May Equivalence and Greenlees-May type duality holds in the settings of P -semiprime and P -semisecond modules. In this section the modules under considerations are bimodules and R -Mod:= R - R -Mod.

We denote by $(R\text{-Mod})_{P\text{-ss}}$ (resp. $(R\text{-Mod})_{P\text{-sp}}$) the subcategory of R -Mod consisting of P -semisecond (resp. P -semiprime) R -modules. A left R -module is said to be P -torsion (resp. P -complete) if and only if $\Gamma_P(M) = M$ (resp. $\Lambda_P(M) = M$).

Proposition 3.1. If M is P -semisecond module and N an R -module, then $\text{Hom}_R(M, N)$ is P -semiprime.

Proof. Suppose that $P^2M = PM$, then $M/P^2M = M/PM$. So, $\text{Hom}_R(R/P^2, \text{Hom}_R(M, N)) \cong \text{Hom}_R(R/P^2 \otimes M, N) \cong \text{Hom}_R(R/P \otimes M, N) \cong \text{Hom}_R(R/P, \text{Hom}_R(M, N))$, then by Proposition 2.5 $\text{Hom}_R(M, N)$ is P -semiprime. For the converse assume that $\text{Hom}_R(M, N)$ is P -semiprime module. Then, $\text{Hom}_R(R/P^2, \text{Hom}_R(M, N)) \cong \text{Hom}_R(R/P, \text{Hom}_R(M, N))$.

$\text{Hom}_R(R/P^2 \otimes M, N) \cong \text{Hom}_R(R/P \otimes M, N)$ if and only if $\text{Hom}_R(M/P^2M \otimes N) \cong \text{Hom}_R(M/PM, N)$. Since N reflects isomorphism, $M/P^2M \cong M/PM$. So, M is P -semisecond. \square

Proposition 3.2. For any ideal P of a ring R we have:

1. $R/P \otimes -$ and $\text{Hom}_R(R/P, -)$ are idempotent functors from R -Mod to $(R\text{-Mod})_{P\text{-ss}} \cap (R\text{-Mod})_{P\text{-sp}}$.
2. For any R -module M , $R/P \otimes \text{Hom}_R(R/P, M) \cong \text{Hom}_R(R/P, M)$ and $\text{Hom}_R(R/P, R/P \otimes M) \cong R/P \otimes M$.
3. For any R -module M , the R -modules $\text{Hom}_R(R/P, M)$ and $R/P \otimes M$ are both P -torsion and P -complete.

Proof.

1. $R/P \otimes (R/P \otimes M) \cong (R/P \otimes R/P) \otimes M \cong R/P \otimes M$ and $\text{Hom}_R(R/P, \text{Hom}_R(R/P, M)) \cong \text{Hom}_R(R/P \otimes R/P, M) \cong \text{Hom}_R(R/P, M)$. For any

R -module M , $R/P \otimes M \cong M/PM$ and $\text{Hom}_R(R/P, M) \cong (0 :_M P)$. Also, $P(M/PM) = 0$ and $P(0 :_M P) = 0$. So, the R -modules M/PM and $(0 :_M P)$ are P -semisecund. It is also easy to see that $(0 :_{(0 :_M P)} P) = (0 :_{(0 :_M P)} P^2) = (0 :_M P)$ and $(\bar{0} :_{M/PM} P) = (\bar{0} :_{M/PM} P^2) = M/PM$. Thus, the R -modules M/PM and $(0 :_M P)$ are P -semiprime.

2. $R/P \otimes \text{Hom}_R(R/P, M) \cong \frac{\text{Hom}_R(R/P, M)}{P\text{Hom}_R(R/P, M)} = \text{Hom}_R(R/P, M)$. $\text{Hom}_R(R/P, R/P \otimes M) \cong \text{Hom}_R(R/P, M/PM) = (\bar{0} :_{M/PM} P) = M/PM \cong R/P \otimes M$.

3. The following maps hold true:

- a. $\text{Hom}_R(R/P, \text{Hom}_R(R/P, M)) \hookrightarrow \Gamma_P(\text{Hom}_R(R/P, M)) \hookrightarrow \text{Hom}_R(R/P, M)$.
- b. $\text{Hom}_R(R/P, R/P \otimes M) \hookrightarrow \Gamma_P(R/P \otimes M) \hookrightarrow R/P \otimes M$.
- c. $\text{Hom}_R(R/P, M) \twoheadrightarrow \Lambda_P(\text{Hom}_R(R/P, M)) \twoheadrightarrow R/P \otimes \text{Hom}_R(R/P, M)$.
- d. $R/P \otimes M \twoheadrightarrow \Lambda_I(R/P \otimes M) \twoheadrightarrow R/P \otimes (R/P \otimes M)$,

where \hookrightarrow denotes a monomorphism and \twoheadrightarrow denotes epimorphism. The first and the last maps are all isomorphisms since $\text{Hom}_R(R/P, M)$ and $R/P \otimes M$ are idempotent. Moreover, $\text{Hom}_R(R/P, M)$ and $R/P \otimes M$ are P -torsion and P -complete. Invariance of $R/P \otimes M$ and $\text{Hom}_R(R/P, M)$ under the functor $\text{Hom}_R(R/P, -)$ and $R/P \otimes -$ respectively shows that the morphisms in (b) and (c) maps are all isomorphisms. This shows that $R/P \otimes M$ and $\text{Hom}_R(R/P, M)$ are P -torsion and P -complete respectively. □

3.1. Greenless-May type Duality. In general, functors Γ_P and Λ_P are not adjoint to each other. However, over commutative ring their derived functors $\mathbf{R}\Gamma_P$ and $\mathbf{L}\Lambda_P$ on the derived category of R -modules are adjoint this is what is called Greenless-May duality, see [13, Theorem 7.12]. In this subsection we show that the functors Γ_P and Λ_P are adjoint in the category of P -semiprime modules and P -semisecund modules, for an ideal P of R (noncommutative).

Lemma 3.3. For any ideal P of a ring R ,

1. The functor $\Gamma_P(-) : (R\text{-Mod})_{P\text{-sp}} \rightarrow (R\text{-Mod})_{P\text{-ss}}$ is idempotent and for any $M \in (R\text{-Mod})_{P\text{-sp}}$, $\Gamma_P(M) \cong \text{Hom}_R(R/P, M)$.
2. The functor $\Lambda_P : (R\text{-Mod})_{P\text{-ss}} \rightarrow (R\text{-Mod})_{P\text{-sp}}$ is idempotent and for any $M \in (R\text{-Mod})_{P\text{-ss}}$, $\Lambda_P(M) \cong R/P \otimes_R M$.

Proof. 1. It follows from Proposition 2.5 (5) and Proposition 3.2 (1).

2. Follows from Proposition 2.10 (3) and Proposition 3.2 (1). \square

Theorem 3.4 (GM type Duality in $R\text{-Mod}$). For any ideal P of a ring R and any $N \in (R\text{-Mod})_{P\text{-sp}}$ and $M \in (R\text{-Mod})_{P\text{-ss}}$,

$$\text{Hom}_R(\Lambda_P(M), N) \cong \text{Hom}_R(M, \Gamma_P(N)).$$

Proof. Consider the functor $\Gamma_P(-) : (R\text{-Mod})_{P\text{-sp}} \rightarrow (R\text{-Mod})_{P\text{-ss}}$. For any module $M \in (R\text{-Mod})_{P\text{-sp}}$, $\Gamma_P(M) \cong \text{Hom}_R(R/P, M)$, Lemma 3.3 (1). However, the functor $R/P \otimes -$ is left-adjoint to $\text{Hom}_R(R/P, -)$. By uniqueness of adjoints, the functor $\Lambda_P(-) : (R\text{-Mod})_{P\text{-ss}} \rightarrow (R\text{-Mod})_{P\text{-sp}}$ which has the property that for all $M \in (R\text{-Mod})_{P\text{-ss}}$ $\Lambda_P(M) \cong R/P \otimes M$, Lemma 3.3 (2). Then, Λ_P is the left adjoint of Γ_P . \square

3.2. Matlis-Greenless-May Equality. Let R be a commutative ring and $\mathbf{D}(R)$ denote the derived category of the abelian category $R\text{-Mod}$. The Matlis-Greenless-May Equivalence (MGM) duality on derived category is given as:

Theorem 3.5. [MGM Equivalence] [13, Theorem 7.11] Let R be a ring, and P be a weakly proregular ideal in it.

1. If $M \in \mathbf{D}(R)$, then $\mathbf{R}\Gamma_P(M) \in \mathbf{D}(R)_{P\text{-tor}}$ and $\mathbf{L}\Lambda_P(M) \in \mathbf{D}(R)_{P\text{-com}}$.
2. The functor $\mathbf{R}\Gamma_P(-) : \mathbf{D}(R)_{P\text{-com}} \rightarrow \mathbf{D}(R)_{P\text{-tor}}$ is an equivalence, with quasi-inverse $\mathbf{L}\Lambda_P$.

In this subsection we prove the MGM Equality in the setting of P -semiprime and P -semisecund modules.

Proposition 3.6. A left R -module M is P -torsion and P -semiprime if and only if M is P -complete and P -semisecund.

Proof. Suppose M be P -torsion and P -semiprime. $M = \Gamma_P(M) = \text{Hom}_R(R/P, M) = (0 :_M P)$, hence it follows that $PM = 0$ which implies $P^k M = 0$ for any $k \in \mathbb{Z}^+$ then $\Lambda_P(M) = M$ and $P^2 M = PM$. Conversely, let M be an P -complete and P -semisecund. To show it is P -semiprime, let $P^k M = 0$ for some $k \in \mathbb{Z}^+$, but since M is P -complete the previous relation satisfied for all $k \in \mathbb{Z}^+$, thus $PM = 0$. Now by Proposition 2.6, $\Gamma_P(M) = \text{Hom}_R(R/P, M) = (0 :_M P) = M$. \square

Lemma 3.7. If P is an ideal of a ring R and M a P -semiprime (resp. P -semisecund) R -module, then $\Gamma_P(M)$ (resp. $\Lambda_P(M)$) is a P -complete (resp. P -torsion) R -module.

Proof. Suppose M is P -semiprime. Then by Proposition 3.2 $\Gamma_P(M) = \text{Hom}_R(R/P, M)$

is both an P -semiprime and P -semisecund R -module. To show that $\Gamma_P(M)$ is P -complete, $\Lambda_P(\Gamma_P(M)) \cong R/P \otimes \text{Hom}_R(R/P, M) \cong \text{Hom}_R(R/P, M) \cong \Gamma_P(M)$. Let M be P -semisecund, by Proposition 3.2, $\Lambda_P(M) \cong R/P \otimes M$ which is also both an P -semiprime and P -semisecund R -module. Now, $\text{Hom}_R(R/P, \Lambda_P(M)) \cong \text{Hom}_R(R/P, R/P \otimes M) \cong R/P \otimes M \cong \Lambda_P(M)$. This proves that $\Lambda_P(M)$ is P -torsion. \square

Let $\mathcal{C} := (R\text{-Mod})_{P\text{-com}} \cap (R\text{-Mod})_{P\text{-ss}}$ and $\mathcal{T} := (R\text{-Mod})_{P\text{-tor}} \cap (R\text{-Mod})_{P\text{-sm}}$.

Theorem 3.8 (MGM Equality). Let P be any ideal of a ring R ,

1. If $M \in (R\text{-Mod})_{P\text{-sm}}$, then $\Gamma_P(M) \in \mathcal{C}$ and if $M \in (R\text{-Mod})_{P\text{-ss}}$, then $\Lambda_P(M) \in \mathcal{T}$.
2. The functor $\Gamma_P(-) : (R\text{-Mod})_{P\text{-sm}} \rightarrow (R\text{-Mod})_{P\text{-ss}}$ restricted to \mathcal{T} is equality between \mathcal{C} and \mathcal{T} with quasi inverse Λ_P .

Proof. 1. Let $M \in (R\text{-Mod})_{P\text{-sm}}$ then by Theorem 3.4 it follows that $\Gamma_P(M) \in (R\text{-Mod})_{P\text{-ss}}$ and by Lemma 3.7, $\Gamma_P(M) \in (R\text{-Mod})_{P\text{-com}}$. Then $\Gamma_P(M) \in \mathcal{C}$. Similarly, applying Theorem 3.4 and Lemma 3.7 we get $\Lambda_P(M) \in \mathcal{T}$.

2. Proposition 3.6 and Lemma 3.7 assures that there is equality between the categories \mathcal{C} and \mathcal{T} which is the Matlis-Greenless-May Equality for R -modules holds. \square

4. THE FUNCTOR $P\Gamma_P$ OVER RINGS

In this section we study some properties of the ideal $P\Gamma_P(R)$ and we use it to formulate Köthe conjecture.

Lemma 4.1. 1. If P is a left ideal of R , then $P\Gamma_P(R)$ is a two sided ideal of R .

2. If P is a right ideal of R , then $P\Gamma_P(R)$ is a right ideal of R .

Proof. 1. Suppose P is a left ideal of R . Let $r \in R$ and $x \in P\Gamma_P(R)$, thus $x = \sum_{i=1}^l a_i r_i$ such that $P^{k_i} r_i = 0$, for some positive integers k_i , where $a_i \in P$ and $r_i \in R$. Now, $rx = \sum_{i=1}^l (ra_i)r_i$,

since P is left ideal $ra_i \in I$ and by hypothesis $P^{k_i}r_i = 0$. Then $x \in P\Gamma_P(R)$ and hence $P\Gamma_P(R)$ is left ideal. To show it is right ideal, $xr = \sum_{i=1}^l a_i(r_i r)$, multiplying $P^{k_i}r_i = 0$ from the right by r we get $P^{k_i}r_i r = 0$ which shows that $xr \in P\Gamma_P(R)$ and hence it is right ideal of R .

2. Suppose P is a right ideal of R . Let $r \in R$ and $x \in P\Gamma_P(R)$, thus $x = \sum_{i=1}^l a_i r_i$ such that $P^{k_i}r_i = 0$, for some positive integers k_i , where $a_i \in P$ and $r_i \in R$. To show $P\Gamma_P(R)$ is right ideal of R , $xr = \sum_{i=1}^l a_i(r_i r)$, multiplying $P^{k_i}r_i = 0$ from the right by r we get $P^{k_i}r_i r = 0$ which shows that $xr \in P\Gamma_P(R)$ and hence it is right ideal of R . □

Proposition 4.2. For any right ideal P of a ring R , $P\Gamma_P(R)$ is a nil right ideal.

Proof. By Lemma 4.1 (2), $P\Gamma_P(R)$ is a right ideal, whenever P is a right ideal of R . Let $y \in P\Gamma_P(R)$. Then $y = \sum_{i=1}^l a_i r_i$ where $r_i \in R$ and $a_i \in P$ such that $P^{n_i}r_i = 0$ for each $1 \leq i \leq l$ and let $n = n_1 + \dots + n_l$. Then, $y^n = (\sum_{i=1}^l a_i r_i)^n = (a_1 r_1)^n + (a_1 r_1)(a_2 r_2)^{n-1} + \dots + (a_1 r_1)(a_i r_i)^{n-1} + \dots$ (l^n terms) each of total degree n . Now, $(a_1 r_1)(a_i r_i)^{n-1} = (a_1 r_1)(a_i r_i)(a_i r_i)^{n-2} \dots (a_i r_i)$, i.e., this is a product of one $a_1 r_1$ term, $n-2$ terms of $a_i r_i$, one term of a_i and r_i , then $(a_1 r_1)(a_i r_i)^{n-1} = (a_1 r_1)(a_i r_i)(a_i r_i)^{n-2} \dots (a_i r_i) \subseteq P P^{n-2} P r_i = P^n r_i = 0$. In a similar fashion every term in the expression y^n undergoes such steps and then $y^n = 0$ and hence $P\Gamma_P(R)$ is a nil right ideal. □

Corollary 4.3. Let R be a ring and \mathcal{U} denotes the upper nilradical of R .

1. For any ideal P of R
 - a. $P\Gamma_P(R)$ is nil.
 - b. $\sum_{P \triangleleft R} P\Gamma_P(R) \subseteq \mathcal{U}(R)$.
2. If R is P -torsion for any ideal P of R , then every ideal P is nil and $\sum_{P \triangleleft R} P\Gamma_P(R) = \mathcal{U}(R)$

Proof. 1. a and b are direct consequences of Proposition 4.2.
 2. By hypothesis $\Gamma_P(R) = R$ for all ideals I of R which implies $P\Gamma_P(R) = PR = P$. By 1a) P is nil, so $\sum_{P \triangleleft R} P\Gamma_P(R) = \sum_{P \triangleleft R} P = \mathcal{U}(R)$. □

Corollary 4.4. *Let R be a right noetherian ring and P be a right ideal of R , then $P\Gamma_P(R)$ is nilpotent.*

Proof. By proposition 2.9, $P\Gamma_P(R)$ is a nil right ideal, for any right ideal P of R then by [10, Levitzki's Theorem] $P\Gamma_P(R)$ is nilpotent. \square

Proposition 4.5. For a ring R , $P\Gamma_P(R[x]) = (P\Gamma_P(R))[x]$.

Proof. Let $f(x) \in P\Gamma_P(R)[x]$. Then $f(x) = \sum_{i=0}^n a_{ij}x^i$, where $a_{ij} \in P\Gamma_P(R)$ which implies for each i there exists S_{ij} and $k_i \in \mathbb{Z}^+$ such that $I^{k_i}S_{ij} = 0$ and $a_{ij} = \sum_{j=0}^k r_{ij}s_{ij}$. Now, $f(x) = \sum_{i=0}^n a_{ij}x^i = \sum_{i=0}^n (\sum_{j=0}^k r_{ij}s_{ij})x^i = \sum_{i=0}^n (r_{i0}s_{i0} + \cdots + r_{ik}s_{ik})x^i = \sum_{i=0}^n r_{i0}(s_{i0}x^i) + \cdots + r_{ik}(s_{ik}x^i)$. Since $I^{k_i}S_{ij} = 0$ it follows that $s_{ij}x^i \in \Gamma_P(R[x])$. Thus, $f(x) \in P\Gamma_P(R[x])$. Suppose $g(x) = \sum_{i=0}^n a_i f_i(x) \in P\Gamma_P(R[x])$, where $f_i(x) = (r_0 + r_1x + \cdots + r_nx^n)_i$ and $P^{k_i}f_i(x) = 0$ which implies $P^{k_i}r_i = 0$ for each $i, 0 \leq i \leq n$ and thus $\sum_{i=0}^n a_i r_i \in P\Gamma_P(R)$. Therefore, $g(x) = \sum_{i=0}^n a_i f_i(x) = \sum_{i=0}^n (a_i r_i)x^i$, so $g(x) \in P\Gamma_P(R)[x]$. \square

Proposition 4.6. Let J_1 and J_2 be ideals of R and P be a right ideal of R . Then $P\Gamma_P(J_1) + P\Gamma_P(J_2)$ is a nil right ideal of R .

Proof. $P\Gamma_P(J_1) + P\Gamma_P(J_2) = P\Gamma_P(J_1 + J_2) \subseteq P\Gamma_P(R)$. From Proposition 4.2 $P\Gamma_P(R)$ is a right nil ideal of R , then it follows that $P\Gamma_P(J_1) + P\Gamma_P(J_2)$ is a right nil ideal of R . \square

Corollary 4.7. If R is a ring such that all its right sided nil ideals are of the form $P\Gamma_P(J)$ for some ideal J of R , then R satisfies the Köthe conjecture

question 4.8. Let P_1 and P_2 be two right ideals of a ring R . Is the sum $P_1\Gamma_{P_1}(R) + P_2\Gamma_{P_2}(R)$ a nil right ideal? A negative answer to this question would answer the Köthe conjecture in the negative.

question 4.9. Is $\sum_{P \triangleleft_r R} P\Gamma_P(R) = \mathcal{U}(R)$? A positive answer to this question would solve the Köthe conjecture in the affirmative.

Acknowledgments

The authors acknowledge support from the International Science Program (ISP), EMS-Simons for Africa and the Eastern Africa Algebra Research Group (EAALG). The authors are grateful to Professor David Ssevviiri for comments.

REFERENCES

- [1] S. Agata, On some results related to Köthe's conjecture, *Serdica Math. J.*, **27** (2) (2001), 159–170.
- [2] R. Beyranvand and F. Rastgoo, Weakly second modules over noncommutative rings, *Hacetatepe J. Math. Stat.*, **45** (5) (2016), 1355–1366.
- [3] M. P. Brodmann and R. Y. Sharp, *Local cohomology: an algebraic introduction with geometric applications*, Cambridge univ. press **136** (2012).
- [4] T. Cheatham and E. Enochs, Injective hulls of flat modules, *Comm. Algebra*, **8**(20) (1980), 1989–1995.
- [5] D. J. Fieldhouse, Pure theories, *Math. Ann.* **184** (1969), 118.
- [6] J. García and J. M. Hernández, When is the category of flat modules abelian?, *Fundam. Math.* **147** (1) (1995), 83–91.
- [7] S. Jøndrup and D. Simson, Indecomposable modules over semiperfect rings, *J. Algebra*, **73** (1) (1981), 23–29.
- [8] G. Köthe. Die Struktur der Ringe, deren Restklassenring nach dem Radikal vollständig irreduzibel ist. *Math. Z.* **32** (1930), 161186.
- [9] A. Kyomuhangi and D. Ssevviiri, The locally nilradical for modules over commutative rings, *Beitr. Algebra Geom.* **61**(4) (2020), 759–769.
- [10] T. Y. Lam, *A first course in noncommutative rings*, **131** (1991), Springer.
- [11] T. K. Lee and Y. Zhou, Reduced modules, rings, modules, algebras and abelian groups, *Lecture Notes in Pure and Appl. Math.* **236** (2004), 365–377.
- [12] L. Némec, T. Bican, P. Kepka, Rings, modules and preradicals, *Lect. notes in pure and appl. math.* **75** (1982).
- [13] M. Porta, L. Shaul and A. Yekutieli, On the homology of completion and torsion, *Algebr. Represent. Theory*, **17**(1) (2014), 31-67.
- [14] E. R. Puczyłowski, Questions related to Köethe's nil ideal problem, *Algebra and its applications, contemporary mathematics*, **419** (2006), 269–283.
- [15] S. P. Redmond, An ideal-based zero-divisor graph of a commutative ring, *Commun. Algebra*, **31**(9) (2003), 4425–4443.
- [16] M. B. Rege and A. M. Buhphang, On reduced modules and rings, *Int. Elect. J. Algebra* **3** (2008), 58–74.
- [17] J. J. Rotman, *An introduction to homological algebra*, **2** (2009), Springer.
- [18] D. Ssevviiri Applications of reduced and coreduced modules II, arXiv preprint arXiv:2205.13241, (2023).
- [19] D. Ssevviiri, Applications of reduced and coreduced modules I, *Int. Electron. J. Algebra* (2023), DOI: 10.24330/ieja.1299587.
- [20] H. Tachikawa, QF-3 rings and categories of projective modules, *J. Algebra*, **28** (1974), 408–413.
- [21] D. Ssevviiri and N. Groenewald, Generalization of nilpotency of ring elements to module elements, *Commun. Algebra*, **42** (2) (2014), 571–577.

Teklemichael Worku Bihonegn

Department of Mathematics, Addis Ababa University , P.O.Box 1176, Addis Ababa, Ethiopia

Email: teklemichael.worku@aau.edu.et

Tilahun Abebaw

Department of Mathematics, Addis Ababa University, P.O.Box 1176, Addis Ababa,
Ethiopia

Email: tilahun.abebaw@aau.edu.et

Nega Arega

Department of Mathematics and Statistics, The Namibia University of Science and
Technology, Namibia

Email: nechere@nust.na